

STUDIES IN CYCLOADDITION REACTIONS : REGIOSPECIFIC CYCLOADDITION
REACTION OF NITRILE IMINES WITH 2H-1-BENZOPYRAN-2-THIONE


Arpan Kumar Baruah, Dipak Prajapati, and Jagir Singh Sandhu*
Division of Drugs and Pharmaceutical Chemistry
Regional Research Laboratory, Jorhat 785006, India

Abstract - Nitrile imines (1) reacted with 2H-1-benzopyran-2-thione to afford cycloadducts (3) in good yields. There was no evidence for the formation of any other regioisomer or the reaction occurring at the thione site.

Cycloaddition chemistry have made considerable impact in organic synthesis¹ and in recent years there has been several reports describing synthesis of complex natural products using this chemistry². The initial recognition of various dipoles³ and dipolarophiles aroused tremendous interest in this area of research and rationalization of mechanistic aspects⁴. During the past several years we have been studying the behaviour of various dipoles towards substrates having more than one dipolarophilic site. In these investigations when 1-aza-1,3-butadienes were reacted with dipoles like benzonitrile oxides⁵ nitrilimines⁶ and aziridines⁷ only more polar carbon-nitrogen double bond reacted and carbon-carbon double bond remained intact. In the present investigation we report our preliminary results on another bifunctional substrate, 2H-1-benzopyran-2-thione which reacted preferentially with nitrilimines at carbon-carbon double bond leaving the thione site intact⁸.

The thione (2) and hydrazinoyl halide (1a) were reacted in equimolar proportions in dry chloroform in the presence of dry triethyl amine at 0-5°C and the reaction mixture was stirred for 30 min. The removal of solvent gave a residue which was dissolved in dry benzene and triethylamine hydrobromide thus precipitated was filtered and solvent was removed under vacuum. The residue thus obtained was further purified by column chromatography, and the compound obtained was essentially a single product (3a) (vide tlc) which was readily

crystalizable from light petroleum (bp 60-80°C) in 70% yield as yellow crystals mp 135-136°C. $^1\text{H NMR}$ (60 MHz, CDCl_3) δ : 2.40 (s, 3H), 5.5 (d,



	R	R'	X
a	CH_3CO	Br	Br
b	CH_3CO	CH_3	Br
c	$\text{C}_2\text{H}_5\text{COO}$	Br	Br
d	$\text{C}_2\text{H}_5\text{COO}$	CH_3	Br
e	C_6H_5	H	Cl.

1H, $J=11\text{Hz}$), 6.8-7.1 (complex multiplet, 9H) ; ν_{max} (KBr), 1250 (lactone, C-O-C), 1505 (C=S), 1695 (C=O). The mass spectrum showed a strong molecular ion at m/z 401. Similarly were obtained compounds (3b-e) in 70-75% yield and their characteristics are recorded in the Table.

It was remarkable to note that the cycloadducts (3a-e) could also be obtained in excellent yields by reacting (1) with (2) under two phase system using TEBA (Tetrabutylammonium bromide) as a phase transfer catalyst and 25% sodium bicarbonate as a base and following rest of the procedure similar to the one as described above. Apparently this procedure involves the slow production of the dipole and eliminates any formation of dimeric products of dipoles.

Table : Characteristic data of cycloadducts (3a-e)*

Compd	Yield (%)	Yield in TPS (%)	Mp °C	IR (KBr) cm ⁻¹	¹ H NMR (60MHz, CDCl ₃) δ ppm.	MS (M ⁺) m/z
3a	70	90	135-136	1250, 1505, 1605, 1695.	2.4(s, 3H), 5.5(d, 1H, J=11Hz), 6.8-7.1(m, 9H).	401, 369, 358, 245.
3b	75	92	110-112	1250, 1510, 1605, 1690.	2.2(s, 3H), 2.4(s, 3H), 5.5(d, 1H, J=11Hz), 6.1-7.05(m, 9H).	336, 304, 293, 245.
3c	72	91	103-104	1250, 1490, 1610, 1720.	1.1(t, 3H), 3.9(q, 2H), 5.6(d, 1H, J=11Hz), 6.3-7.0(m, 9H).	431, 399, 358, 275.
3d	75	90	92-94	1245, 1490, 1610, 1720.	1.2(t, 3H), 4.0(q, 2H), 5.5(d, 1H, J=11Hz), 6.2-7.0(m, 9H).	366, 334, 293, 275.
3e	70	88	113-114	1225, 1500, 1605.	5.5(d, 1H, J=11Hz), 6.15-7.30(m, 15H).	356, 324, 279, 265.

* All the compounds reported here gave satisfactory elemental analyses.

ACKNOWLEDGEMENT

We thank the Analytical Chemistry Division of this Laboratory for some of the spectral analyses. One of us (A.K.B.) also thanks the Council of Scientific and Industrial Research (CSIR) New Delhi for the award of a research fellowship.

REFERENCES

1. For recent review see :
 - a) A. Padwa, Angew. Chem., 1967, 88, 131 ; Ibid ; Int. Ed. Engl., 1976, 15, 123.
 - b) W. Oppolzer, Angew. Chem. Int. Ed. Engl., 1977, 16, 10.
 - c) R. Huisgen, Angew. Chem. Int. Ed. Engl., 1980, 19, 966.
 - d) D.L. Boger, Tetrahedron, 1983, 39, 2869.
2. a) J.J. Tufariello, H. Meckler, K. Pushpananda and A. Senaratne, Tetrahedron, 1985, 41, 3447.
 - b) A.E. Walts and W.R. Roush, Tetrahedron, 1985, 41, 3463.
3. a) R. Huisgen, Angew. Chem., 1963, 75, 604, 742.
 - b) R. Huisgen, Angew. Chem., Int. Ed. Engl., 1963, 2, 565, 633.
4. a) R. Huisgen, J. Org. Chem., 1968, 33, 2291 ; Ibid, 1976, 41, 403 ; Pure Appl. Chem., 1981, 53, 171.
 - b) R.A. Firestone, J. Org. Chem., 1968, 33, 2285 ; J. Chem. Soc. A., 1970, 1570 ; J. Org. Chem., 1972, 37, 2181 ; Tetrahedron, 1977, 33, 3009.
 - c) R.D. Harcourt and R.D. Little, J. Am. Chem. Soc., 1984, 106, 41 and references cited therein.
5. N. Singh and J.S. Sandhu, Tetrahedron Lett., 1968, 42, 4453.
6. a) N. Singh, J.S. Sandhu and S. Mohan, J. Chem. Soc. Chem. Commun., 1969, 387.
 - b) D. Prajapati, J.S. Sandhu, T. Kametani, H. Nagase, K. Kawai and T. Honda, Heterocycles, 1985, 23, 1123.
 - c) D. Prajapati, J.S. Sandhu and J.N. Baruah, J. Chem. Research (S), 1984, 56.
 - d) D. Prajapati and J.S. Sandhu, Heterocycles, 1985, 23, 1143.
7. B. Sain and J.S. Sandhu, Ind. J. Chem., 1985, 24B, 291.
8. Thione is a good dipolarophile e.g. see :
 - a) T. Katada, S. Eguchi and T. Sasaki, J. Chem. Soc. Perkin Trans. I, 1984, 2641.
 - b) R. Huisgen, C. Fulka, I. Kalwisch, L. Xingya, G. Mloston, J.R. Moran and A. Probstl, Bull. Soc. Chims. Belg., 1984, 93, 511.
 - c) K. Bunge, R. Huisgen, R. Raab and H.J. Sturm, Chem. Ber., 1972, 105, 1307.
 - d) R. Huisgen, G. Mloston and A. Probstl, Tetrahedron Lett., 1985, 26, 4431 and references cited therein.

Received, 14th January, 1986