

INTRAMOLECULAR CYCLISATION USING METHYL(BISMETHYLTHIO)SULPHONIUM SALTS.
PART 5.¹ SYNTHESIS OF FUNCTIONALIZED 4,5-DIHYDRO-3,1-BENZOXAZEPINES

Giuseppe Capozzi*

Centro di Studio CNR sulla Chimica e La Struttura dei Composti Eterociclici, Dipartimento di Chimica Organica 'Ugo Schiff' - Università di Firenze - 50100 Firenze - Italy

Rosaria Ottana¹ and Giovanni Romeo*

Dipartimento Farmaco-Chimico - Università di Messina - 98100 Messina - Italy


Abstract - The reaction of methyl(bismethylthio)sulphonium hexachloroantimonate **1** with N-(2-allylphenyl)benzamide **2** gives the 4-methylthio-methyl-4,5-dihydro-3,1-benzoxazepine **4**. Other electrophilic reagents reacts with **2** to give exclusively or preferentially products of addition to the allylic double bond.

In recent years we have introduced the use of methyl(bismethylthio)sulphonium hexachloroantimonate **1** as a synthon of methylsulphenylium ion (MeS^+) for the cyclofunctionalization of properly substituted alkenes and alkynes.

In this way, methylthio-functionalized dihydrobenzofurans,² methylene-oxazoles,³ dihydrooxazines,⁴ indoles and dihydroindoles¹ have been obtained from o-allylphenols, propargylamides, o-vinylbenzanilides, o-vinylbenzenesulphonamides, and o-allylbenzenesulphonamides respectively.

We report here some preliminary results on the application of this reaction to the synthesis of the 3,1-benzoxazepine system for which the available synthetic approaches are limited.⁵

- The suitably substituted alkene **2**, prepared from the corresponding aniline derivative,⁶ was reacted with the sulphonium salt **1** in dichloromethane at 0°C. Addition of n-pentane to the reaction mixture gives the benzoxazepinium hexachloroantimonate **3** which was recrystallized from dichloromethane-n-pentane to give pure **3** in 60% yield⁷ (equation).




Treatment of **3** with aqueous sodium bicarbonate solution and column chromatography (eluant light petroleum - diethyl ether 7:3) gives 4-methylthiomethyl-4,5-dihydro-3,1-benzoxazepine **4** in quantitative yield.

Compound **4**⁸ gave correct elemental analysis and spectroscopic properties in agreement with the proposed structure. In particular the mass spectrum showed the molecular ion at 283 m/z and the ion at 222 m/z, corresponding to the loss of the CH₂SMe moiety as relatively intense peak. The ¹H nmr spectrum showed, besides the S-methyl and aromatic protons resonances, two ABX systems due to the two methylene groups coupled with the same methine proton.⁹ The ¹³C nmr spectrum gave more compelling informations on the structure of **4**:¹⁰ the 4-C was found as doublet at δ 81.71 and the 2-C as a singlet at δ 151.34.

Other electrophilic reagents were also tested for the ring closure of **2**.

The attempt to cyclize **2** by acid catalysis using methanesulphonic acid gives **5** only,¹¹ the addition product of the acid to the carbon-carbon double bond. The reaction of **2** with methanesulphenyl chloride gives a 1:2 mixture of **4** and **6**¹² which is the adduct of the sulphenyl chloride to **2**.



The results above reported clearly show the stronger tendency of the sulphonium salt **1** with respect to other electrophilic reagents to promote the ring closure of **2** to the seven-membered ring product.

This behaviour is possibly due to the absence in this reagent of a strong nucleophilic counterpart which may compete with the poorly nucleophilic amidic oxygen of **2** for the attack at the positively charged intermediate which, in the case of the sulphenic electrophiles, should have the structure of a thiiranium ion.¹³

ACKNOWLEDGMENT

We acknowledge financial support from 'Progetto Finalizzato Chimica Fine e Secondaria', C.N.R.

REFERENCES AND NOTES

1. Part 4, G. Capozzi, R. Ottana', and G. Romeo, Heterocycles, 1986, 24, 583.
2. G. Capozzi, V. Lucchini, F. Marcuzzi, and G. Modena, J. Chem. Soc., Perkin Trans. 1, 1981, 3106.
3. G. Capozzi, C. Caristi, and M. Gattuso, J. Chem. Soc., Perkin Trans. 1, 1984, 255.
4. G. Capozzi, R. Ottana', G. Romeo, and G. Valle, J. Chem. Research (S), 1986, 200
5. J. T. Sharp in "Comprehensive Heterocyclic Chemistry", eds. A. R. Katritzky and C. W. Rees, Pergamon Press, New York, 1984, vol. 7, p. 626; J. K. Landquist in "Comprehensive Organic Chemistry", eds. D. Barton and D. Ollis, Pergamon Press, New York, 1979, vol. 4, p. 1153.
6. Compound 2 was synthesized by reaction of o-allylaniline and benzoyl chloride in the presence of Et₃N in CH₂Cl₂ at room temperature; yield 60%, mp 143-145°C from diethyl ether; ¹H nmr in CDCl₃: δ 8.10-7.10 (m, 10 H, aromatics and NH), 6.06 (m, 1 H), 5.18 (m, 2 H), 3.46 (d, J 5.80 Hz, 2 H). Mass spectrum, molecular ion at m/z 237. Satisfactory elemental analyses were obtained for this compound and for all the other new compounds described hereinafter.
7. mp 99-101°C; ¹H nmr in CD₃COCD₃: δ 8.35-7.47 (m, 9 H, aromatics), 5.65 (m, 1 H) 4.47 (broad s, NH), 3.82 (m, 2 H), 3.30 (m, 2 H), 2.27 (s, 3 H).
8. Compound 4 was purified by column chromatography (SiO₂, eluant light petroleum-diethyl ether, 7:3 v/v) and bulb to bulb distillation at 4 mm Hg, oil bath temperature, 200°C.
9. ¹H nmr in CDCl₃: δ 8.14 (m, 2 H, aromatics), 7.55-7.05 (m, 7 H, aromatics), 4.76 (m, 1 H), 3.21 (m, 2 H), 2.83 (m, 2 H), 2.17 (s, 3 H).
10. ¹³C nmr in CDCl₃: δ 151.34 (s), 143.52 (s), 135.56 (s), 132.45 (s), 130.37 (d), 129.06 (d), 128.93 (d), 128.43 (d), 127.92 (d), 127.51 (d), 124.94 (d), 81.71 (d), 40.05 (t), 39.20 (t), 16.30 (q).
11. The reaction of 2 (1 mmol) with methanesulphonic acid (4 ml) was carried out at room temperature for 30 min. After usual work-up the reaction mixture was

chromatographed (SiO_2 , eluant light petroleum-diethyl ether 7:3 v/v) to give the unreacted **2** (0.4 mmol) and **5** (0.44 mmol, 44% yield) mp 123-125°C. ^1H nmr in CDCl_3 : δ 8.18 (broad s, NH), 7.95-7.15 (m, 9 H, aromatics), 4.90 (m, 1H), 3.10 and 2.92 (AB part of an ABX system, 2H, J_{AB} 14.65 Hz, J_{AX} 4.83 Hz, J_{BX} 8.54 Hz), 2.53 (s, 3H), 1.53 (A part of an A_3X system, 3 H, J_{AX} 6.10 Hz). Mass spectrum, molecular ion at m/z 333.

12. Equimolar amount of **1** and methane sulphenyl chloride in dichloromethane at 0°C were reacted for 2 h giving after work-up and column chromatography (SiO_2 , eluant light petroleum-diethyl ether 7:3 v/v) **4** (25% yield) and **6** (52 % yield). Compound **6** gave correct elemental analysis and a mass spectrum with the molecular ion at 319 m/z . The regiochemistry of the addition of the methanesulphenyl chloride to **1** has been deduced from the chemical shift values of the ^1H nmr spectrum and in particular from the low value of the methine proton (δ 4.25). ^1H nmr in CDCl_3 : δ 8.42 (broad s, NH), 7.95 (m, 2 H, aromatics), 7.42 (m, 7 H, aromatics), 4.25 (m, 1 H), 3.49 and 3.05 (AB part of an ABX system, 2 H, J_{AB} 14.95 Hz, J_{AX} 8.54 Hz, J_{BX} 3.34 Hz), 2.93 (m, 2 H), 2.07 (s, 3 H).

13. For recent reviews, see: G. Capozzi and G. Modena in "Organic Sulfur Chemistry. Theoretical and Experimental Advances", eds. F. Bernardi and A. Mangini, Elsevier, Amsterdam, 1985, p. 246. G. H. Schmid and D. G. Garratt in "The Chemistry of Functional Groups. Supplement A. The Chemistry of Double-Bonded Functional Groups", ed. S. Patai, Wiley, New York, 1977, part 2, p. 828.

Received, 2nd September, 1986