

BIOLOGICAL FORMATION AND CHEMICAL SYNTHESIS OF 2-AMINO-3,6-DIHYDRO-3-METHYL-7H-IMIDAZOLO[4,5-f]QUINOLIN-7-ONE, THE MAJOR METABOLITE OF THE DIETARY CARCINOGEN 2-AMINO-3-METHYL-3H-IMIDAZOLO[4,5-f]QUINOLINE (IQ) BY NORMAL INTESTINAL BACTERIA

Mohammad Bashir and David G. I. Kingston*


Department of Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U.S.A.

Robert J. Carman, Roger L. Van Tassell, and Tracy D. Wilkins


Department of Anaerobic Microbiology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U.S.A.

Abstract—2-Amino-3,6-dihydro-3-methyl-7H-imidazo[4,5-f]quinolin-7-one (2) has been synthesized from 6-bromo-5-nitroquinoline in seven steps. Biological formation of 2 from IQ (1) involves the addition of water from the medium, followed by oxidation.

The formation of various mutagenic heterocyclic amines on pyrolysis of proteinaceous foods has been reported by many investigators,¹ and several reviews of the subject have appeared.² Mutagens of the imidazoquinoline or imidazoquinoxaline class of compounds are the most potent mutagens known in the Ames test, and the parent compound of this group, 2-amino-3-methyl-3H-imidazo[4,5-f]quinoline, or IQ (1) has been shown to be carcinogenic in mice^{1a,3} and in rats.⁴



1



2

Because of the significance of IQ as a possible carcinogen in man, a knowledge of its metabolic fate is important to an assessment of its carcinogenic potential. Some studies have been reported on the fate of IQ in rats, and it was shown that the major urinary mutagen is unchanged IQ, with smaller amounts of possible N-acetyl or 3-N-demethylated products.⁵ It has also been shown that IQ is metabolized to an active N-hydroxy compound by cytochrome P-448.⁶ Recently we reported that IQ is converted to a metabolite identified as 2-amino-3,6-dihydro-3-methyl-7H-imidazo[4,5-f]-quinolin-7-one (HOIQ) (2) by human fecal flora,^{7,8} and we have also shown that this metabolite is a potent direct acting mutagen,⁸ in contrast to IQ which requires activation by liver microsomal enzymes.


Since HOIQ is potentially a greater cancer risk than IQ itself, due to its direct acting nature, it thus became important to prepare it by synthesis, both as a confirmation of structure and to provide adequate material for biological testing. In addition, the mechanism of the unusual formation of an oxidized compound under anaerobic conditions is of interest. In this manuscript, we report the synthesis of HOIQ and a study of its metabolic formation.

RESULTS AND DISCUSSION

The first synthetic route to HOIQ that we attempted was a simple modification of the published route to IQ.⁹ Oxidation of 6-bromo-5-nitroquinoline 3⁹ with m-chloroperbenzoic acid yielded the N-oxide 4. Rearrangement of 4 in the presence of acetic anhydride gave the acetate 5, but reactions of this compound with methylamine, under conditions which were successful when applied to 3,⁹ failed to effect a displacement of bromide. We thus sought an alternate route to avoid this problem.

Reaction of 6-bromo-5-nitroquinoline 3 with methylamine as previously described⁹ gave the aminoquinoline 7, which was acetylated to the acetamidoquinoline 8. Treatment of 8 with m-chloroperbenzoic acid gave the N-oxide 9 in 70% yield. Rearrangement of 9 in the presence of acetic anhydride and sodium acetate, according to a literature procedure,¹⁰ gave substantial amounts of desoxy material (8) in addition to the desired rearrangement product. The desoxy product was presumably formed by attack of acetate ion on the N-acetoxyquinolinium ion to give methyl acetate and carbon dioxide, and the undesired reaction was avoided by the use of acetic anhydride alone for the rearrangement. The 2(1H)-quinolinone 10 was obtained in 50% yield after a work-up which included a mild basic hydrolysis.

Attempted conversion of 10 to the desired desacetyl product 15 produced some mild surprises. Hydrolysis with 30% NaOH resulted in displacement of the N-methylacetamido group by hydroxide ion to give the 6-hydroxy-5-nitro-2(1H)-quinolinone 12. Reduction prior to hydrolysis did not help, since the aminoquinolinone 13 simply underwent cyclization on treatment with base to give the 2,3-



dimethylimidazoquinolin-2-one 14. Vigorous hydrolysis in the presence of hydrochloric acid also yielded an unexpected product in the shape of 5-chloro-6-(methylamino)-2(1H)-quinolinone 11. Similar displacements of nitro groups by chloride ion have been reported when 4-nitroquinoline-N-oxides are heated with hydrochloric acid.¹¹

Conversion of the nitroquinolinone 10 to the desired diaminoquinolinone 16 was finally achieved by mild acid hydrolysis to the aminonitroquinolinone 15, followed by catalytic reduction to 16. Completion of the synthesis was then brought about by the method previously used to make IQ.⁹ The isolated product 2 had identical chemical properties (¹H nmr, tlc, ms) to the metabolite obtained from IQ,⁷ and it also showed the same mutagenicity in the Ames assay.⁸

The biochemical formation of compound 2 under anaerobic conditions is surprising, since formally it is an oxidation product of IQ (1). The hydroxylation of aromatic rings by microorganisms is an important step in the metabolism of various physiologically important compounds, and it occurs by three different pathways. Two of these pathways involve oxidation by dioxygen, with catalysis by monooxygenases or by dioxygenases, and these are the normal pathways for the aerobic metabolism of aromatic compounds.¹² A third pathway involves the addition of water followed by hydride transfer to an electron acceptor, and this is the pathway that has been demonstrated in the oxidation of nicotinic acid under anaerobic conditions.¹³

Incubation of IQ in a culture of the anaerobic bacterium Eubacterium moniliforme VPI 13480, which metabolizes IQ,⁸ with added H₂¹⁸O (20% wt/vol), yielded a sample of HOIQ (2) which contained about 12% ¹⁸O as determined by mass spectrometry. Since a control experiment indicated that oxygen exchange of the amide with water does not occur under the experimental conditions, this result shows that the formation of HOIQ from IQ must involve an initial hydration of the 1,2-bond of the quinoline ring, followed by oxidation of the resulting dihydroquinoline. The electron acceptor for this oxidation step is at present unknown.

In conclusion, we have synthesized the unusual metabolite HOIQ, thus confirming its structure, and have determined that it arises by a hydration-dehydrogenation sequence on the food mutagen IQ. Studies on the biological activity of HOIQ are in progress, and will be reported elsewhere.⁸

EXPERIMENTAL

Melting points (uncorrected) were determined in open capillaries. Proton spectra were obtained in the indicated solvent on an IBM WP-270 nuclear magnetic resonance (nmr) spectrometer; chemical shifts are reported in ppm downfield from internal tetramethylsilane. Splitting patterns are designated as s(singlet), d(doublet), t(triplet), q(quartet), and m(multiplet). Coupling constants are given in Hertz. Infrared spectra (ir) were recorded on a Perkin-Elmer instrument. Ultraviolet (uv) spectra were obtained from solutions in ethanol on a Hitachi-Perkin-Elmer spectrophotometer.

Mass spectra were obtained on a VG 7070 instrument operating in the electron ionization (ei) or chemical ionization (ci) mode. Flash chromatography was carried out on E. Merck silica gel (230-400 mesh) according to the Still procedure.¹⁴ Thin layer chromatography was carried out on E. Merck precoated silica gel plates, 0.2mm. Microanalyses were performed by Micanal (Tucson, Az.)

6-Bromo-5-nitroquinoline-N-oxide(4).

A solution of 3 (1 g, 3.96 mmol) in CHCl_3 (5 ml) was cooled to 0°C and to this solution MCPBA (0.72 g, 4.18 mmol) in 10 ml of CHCl_3 was added dropwise at the same temperature. Stirring was continued for 3 h, and the reaction mixture was kept overnight at 0°C. The resulting solution was washed with saturated NaHCO_3 and then with water. The organic layer was dried (MgSO_4) and evaporated in vacuo. Purification by flash chromatography (CHCl_3) afforded 4 (0.33 g, 31%) which crystallized from EtOH, mp 204-206°C: ^1H nmr (CDCl_3) 7.29-7.51(2H,m), 7.94(1H,d,J=9.3), 8.59(1H,d,J=5.6), 8.77(1H,d,J=9.3); eims m/z (relative intensity) 270,268(M^+ ,100,98), 254,252(30,30), 240,238(36,34), 224,222(42,42), 196,194(25,27), 143(80), 127(90), 115(70); uv λ_{max} 345 nm(ϵ 5913), 236(30107); ir(CHCl_3), 1550, 1360, 1270, 1260, 1160 cm^{-1} .

2-Acetoxy-6-bromo-5-nitroquinoline (5).

Compound 4 (64 mg, 0.238 mmol) was dissolved in 5 ml of acetic anhydride and refluxed for 3 h in an N_2 atmosphere. Solvent was then removed in vacuo. The residue was basified with NH_4OH and extracted with CHCl_3 . The organic layers were combined, dried (Na_2SO_4) and evaporated. Preparative tlc (CHCl_3 : EtOAc, 1:1) yielded 5 (22 mg, 30%) as brown needles from EtOH, mp 83-85°C: ^1H nmr (CDCl_3) 2.43(3H,s), 7.40(1H,d,J=8.9), 7.92(1H,d,J=8.9), 8.01(1H.d.J=9.5), 8.12(1H.d.J=9.5); eims m/z (relative intensity) 312,310(M^+ , 50,44), 270/268 (100,100), 224,222(20,22), 196,194(22,25), 115(50); uv λ_{max} 335nm (ϵ 2713); 325(3488), 310(2713), 275(3100), 232(37209); ir(CHCl_3), 1780, 1540, 1370, 1170 cm^{-1} .

6-Bromo-5-nitro-2(1H)-quinoline(6).

A solution of 5 (22 mg, 0.07 mmol) in EtOH (1.5ml) was refluxed with 40% aq methylamine (0.1 ml) for 1 h. Solvent was removed and the residue was purified by preparative tlc (CHCl_3 : EtOAc, 1:1) to give 6 (18 mg, 94%), mp >290°C(MeOH): ^1H nmr(DMSO) 6.71(1H,d,J=9.9), 7.44(1H,d,J=8.9), 7.66(1H,d,J=9.9), 7.93(1H,d,J=8.9); eims m/z (relative intensity) 270,268 (M^+ ,50,50), 224,222(22,20), 212,210(30,36), 196,194(35,40), 167(20), 115(100); uv λ_{max} 345 nm (ϵ 5376), 275(2150), 238(35483); ir (KBr) 3000, 2900, 1620, 1510, 1460, 1420, 1340, 1220 cm^{-1} .

6-Methylamino-5-nitroquinoline (7).

6-Bromo-5-nitroquinoline (10 g, 39.52 mmol) was heated under reflux in 120 ml of ethanol, and to this refluxing solution 40% aq methylamine (20 g, 258 mmol) was added dropwise over 1 h. After refluxing for another 3 h tlc showed the absence of starting material. The solution was then poured onto ice water and extracted with CHCl_3 . The organic fractions were combined, dried (Na_2SO_4) and evaporated. Crystallization of the residue from acetone gave yellow needles of 7 (6.4 g, 80%), mp 190°C (lit. 194°C); ^1H nmr (CDCl_3) 3.20(3H,d,J=5), 7.33(1H,d,J=9.7), 7.50(1H,dd, $J_1=8.9, J_2=4.2$), 8.09(1H,d,J=9.7), 8.69(1H,dd, $J_1=4.2, J_2=1.4$), 9.15(1H, broad s), 9.18(1H,d,J=8.9); eims m/z (relative intensity) 203(M^+ ,100), 186(12), 173(10), 156(40), 128(60), 115(15), 101(30), 77(20); uv λ_{max} 430 (ϵ 23571), 315(20000), 245(66428); ir(CHCl_3) 3600, 3050, 1620, 1540, 1220, 940 cm^{-1} .

6-(N-Acetyl-N-methylamino)-5-nitroquinoline (8).

Compound 7 (5 g, 24.63 mmol) was dissolved in acetic anhydride (50 ml) and pyridine (10 ml) and refluxed for 8 h in an N_2 atmosphere. Solvent was removed *in vacuo* and the residue was dissolved in CHCl_3 and washed with water. The organic layer was dried (Na_2SO_4) and concentrated to a gum. Purification by flash chromatography (CHCl_3) afforded 8 (5.25 g, 87%) which crystallized from EtOAc, mp 84-85°C: ^1H nmr (CDCl_3) 1.90(3H,s), 3.29(3H,s) 7.27-7.69(2H,m), 8.16(1H,d,J=8.4), 8.38(1H,d,J=9), 9.10(1H,d,J=4.2); cims m/z (relative intensity) 246($\text{M}+1^+$,90), 238(10), 216(20) 204(45), 198(95), 170(30), 145(10); uv λ_{max} 315nm(ϵ 3750), 307(3000), 218(39500); ir(CHCl_3) 3050, 1700, 1560, 1400, 1350, 1200 cm^{-1} . Anal. Calcd for $\text{C}_{12}\text{H}_{11}\text{N}_3\text{O}_3$: C, 58.77; H, 4.48; N, 17.14. Found: C, 58.49; H, 4.15; N, 16.88.

6-(N-Acetyl-N-methylamino)-5-nitroquinoline-N-oxide (9).

A solution of m-chloroperbenzoic acid (4.95 g, 28.77 mmol) in 90ml of CHCl_3 was added dropwise over 2 h, to a stirred solution of 8 (4 g, 16.32 mmol) in 20 ml of CHCl_3 at 0°C in an N_2 atmosphere. The reaction mixture was stirred for an additional 6 h at the same temperature and then kept overnight at 0°C. The solution was then treated with a saturated solution of NaHCO_3 and finally with water. The organic layer was dried (MgSO_4) and evaporated *in vacuo*. Flash chromatography (CHCl_3 , CHCl_3 :MeOH 95:5) gave 9 (2.98 g, 70%) as a yellow solid, which was recrystallized from EtOH, mp 181-182°C: ^1H nmr (CDCl_3) 1.88(3H,s), 3.25(3H,s), 7.50(1H,m), 7.64(2H,m), 8.60(1H,d,J=5.6), 9.0(1H,d,J=9.4); cims m/z (relative intensity) 262($\text{M}+1^+$,90), 246(60), 220(53), 215(100), 199(85), 186(13), 170(20); uv λ_{max} 350 nm(ϵ 7000), 230(27666), 218(28000); ir(CHCl_3) 3050, 1690, 1550, 1360, 1270 cm^{-1} . Anal. calcd. for $\text{C}_{12}\text{H}_{11}\text{N}_3\text{O}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$: C,53.33; H,4.44; N,15.55. Found: C,52.92; H,4.01; N,15.19.

6-(N-Acetyl-N-methylamino)-5-nitroquinolin-2-one (10).

A solution of 9 (2 g, 7.66 mmol) in 50 ml of acetic anhydride was refluxed for 30 h in an N₂ atmosphere. Solvent was then removed *in vacuo* and the residue was basified with NH₄OH and extracted with CHCl₃. The organic phases were combined, dried (Na₂SO₄) and evaporated. Purification by flash chromatography (CHCl₃ → CHCl₃: MeOH, 90:10) afforded 10 (1g, 50%), mp 259-260°C (EtOH): ¹H nmr (CDCl₃) 1.86(3H,s), 3.21(3H,s), 6.90(1H,d,J=9.8), 7.46(1H,d,J=8.8), 7.64(1H,d,J=8.8), 7.74(1H,d,J=9.8), 12.05(1H,bd.s); cims, m/z (relative intensity) 262(M+1⁺,100), 233(30), 220(25), 215(70), 199(12), 186(25), 175(15); uv λ_{max} 335 nm(ε6190), 270(3809), 235(40952); ir(CHCl₃) 3675, 3050, 1690, 1550, 1440, 1240, 1050cm⁻¹. Anal. Calcd. for C₁₂H₁₁N₃O₄: C,55.17; H,4.2; N,16.09. Found: C,54.89; H,4.16; N,16.13.

3,6-Dihydro-2,3-dimethyl-7H-imidazo[4,5-f]quinolin-7-one (14).

A mixture of 10 (50 mg, 0.19 mmol) and 10% Pd/C (10 mg) in 5 ml of acetic acid was hydrogenated at room temperature. After 3 h the catalyst was filtered off and the filtrate was evaporated. The residue was basified with NH₄OH and extracted with CHCl₃, and the CHCl₃ fractions were dried (Na₂SO₄) and evaporated to afford crude product. Purification by preparative tlc (CHCl₃:MeOH, 95:5) afforded the unstable amine 13 (13 mg). Compound 13 (13 mg) was refluxed with 20% NaOH (1ml) for 2 h. After this time water was removed from the reaction mixture and the product was directly purified by preparative tlc to give 14 (4.4 mg, 36.7%): ¹H nmr (CD₃OD) 2.54(3H,s), 3.72(3H,s), 6.60(1H,d,J=9.5), 7.15(1H,d,J=8.8), 7.57(1H,d,J=8.8), 8.40(1H,d,J=9.5); cims, m/z (relative intensity) 214(M+1⁺,100), 199(10), 171(6).

5-Chloro-6-methylaminoquinolin-2-one (11).

A solution of 10 (10 mg, 0.038 mmol) in 5% HCl (1.5ml) was refluxed for 15 h in an N₂ atmosphere. The solution was basified with NH₄OH, extracted with CHCl₃ and the organic phases were combined, dried (Na₂SO₄) and concentrated to dryness *in vacuo*. Purification by preparative tlc (CHCl₃: EtOAc, 1:1) afforded 11 (6 mg, 75%): ¹H nmr(CD₃OD) 2.79(3H,s), 6.55(1H,d,J=9.8), 6.97(1H,d,J=9.0), 7.14(1H,d,J=9.0), 8.09(1H,d,J=9.8); eims m/z (relative intensity) 210,208 (M⁺32,100), 195,193(8.25), 173(5), 166(7).

6-Hydroxy-5-nitro-2(1H)-quinolinone (12).

Compound 10 (80 mg, 0.305 mmol) in 2.5ml of 30% NaOH was refluxed for 4 h in an N₂ atmosphere. Water was removed *in vacuo* and the residue was purified by flash chromatography (EtOAc→EtOAc:MeOH, 70:30) to yield 12 (60 mg, 95%); ¹H nmr (CD₃OD), 6.36(1H,d,J=9.3), 6.70(1H,d,J=9.0), 7.10(1H,d,J=9.3), 7.76(1H,d,J=9.0).

6-Methylamino-5-nitro-2(1H)-quinolinone (15).

Compound 10 (0.7 g, 2.68 mmol) was dissolved in 5% HCl (30 ml) and heated to 70°C for 24 h in an N₂ atmosphere. After this time a red precipitate was filtered off, washed with water and basified with NH₄OH. The basic solution was extracted with a large excess of CHCl₃. The CHCl₃ fractions were combined, washed with water, dried (Na₂SO₄) and concentrated, to give 15 (406 mg, 70%) as a red solid. Recrystallization from acetic acid give red crystals, mp 310-313°C: ¹H nmr (DMSO) 2.92(3H,d,J=4.9), 6.60(1H,d,J=10), 7.30(1H,d,J=9.4), 7.5(1H,d,J=9.4), 7.73(1H,br s), 8.25(1H,d,J=10), 11.89(1H,a); eims, m/z (relative intensity) 219(M⁺,100), 202(15), 185(10), 173(30), 128(15), 116(50), 89(15); uv λ_{max} 470nm (ε5555), 340(2777), 248(27777); ir(KBr) 3500, 2900, 1740, 1700, 1670, 1550, 1400, 1365, 1340cm⁻¹.

5-Amino-6-methylamino-2(1H)-quinolinone (16).

A stirred heterogeneous mixture of 15 (0.32 g, 1.46 mmol) and 10% Pd/C (100 mg) in 100 ml of acetic acid was hydrogenated at room temperature for 10 h. The catalyst was removed by filtration, the filtrate was evaporated, and the residue was basified with NH₄OH and extracted with EtOAc. Organic phases were combined, dried (Na₂SO₄), and evaporated *in vacuo*. Purification by flash chromatography (CHCl₃--->CHCl₃: MeOH, 90:10) yielded 16 (135mg, 49%). ¹H nmr (DMSO) 2.70(3H,d,J=5), 4.09(1H,q,J=5), 5.23(2H,s), 6.21(1H,d,J=9.7), 6.48(1H,d,J=8.5), 6.90(1H,J=8.5), 8.08(1H,d,J=9.7), 11.12(1H,s); eims, m/z (relative intensity) 189(M⁺, 90), 174(98), 147(35), 129(7), 119(10), 104(12), 92(8).

2-Amino-3,6-dihydro-2-3-methyl-7H-imidazo[4,5-f]quinolin-7-one (HOIQ, 2).

Cyanogen bromide (126 mg, 1.18 mmol) and 16 (113 mg, 0.5 mmol) were dissolved in 8 ml of MeOH and stirred overnight at room temperature. Next day the hydrobromide 17 was filtered off, basified with 1N NaOH solution, and filtered. The residue was washed with water and finally with EtOH. Crystallization of the residue from EtOH yielded 2 (60 mg, 46.8%) as light green iridescent crystals, mp>360°C: ¹H nmr (DMSO) 3.50(3H,s), 6.50(1H,d,J=9.5), 6.60(2H,s), 6.51(1H,d,J=8.5), 7.30(1H,d,J=8.5), 8.10(1H,d,J=9.5), 11.48(1H,s); eims, m/z (relative intensity) 214(M⁺, 100), 199(30), 186(15), 171(20); uv λ_{max} 360 nm (ε3333), 330(9444), 315(1111), 270(15555), 225(30555); ir (KBr) 3300, 3100, 1630, 1600, 1580, 1550, 1440, 1380, 1340, 1240, 1220 cm⁻¹.

The isolated product was identical (¹H nmr, tlc, co-tlc) with the metabolite isolated from feces.⁷

Metabolism of IQ in H₂¹⁸O.

Brain heart infusion (BHI) broths (Difco) were prepared using [¹⁸O] water (20 atom % ¹⁸O; MSD isotopes, Montreal, Canada) and supplemented with [¹⁴C] IQ (5 µg/ml). Each tube was inoculated

with 100 μ l of an actively growing BHI culture of Eubacterium moniliforme VPI 13480, a known metabolizer of IQ¹⁶. The tubes were incubated anaerobically for 18 h at 37°C after which HOIQ and the residue of the labelled substrate were extracted with blue cotton.⁷ HOIQ was purified as previously described,⁷ and analyzed by mass spectrometry. HOIQ prepared from H₂¹⁶O had peaks in the molecular ion region at m/z 214 (M⁺, 100), 215(13.6), 216(2.0), while HOIQ prepared from H₂¹⁸O as described had peaks at m/z 214 (M⁺, 100), 215(16.8), and 216(16.7). A control experiment in which HOIQ was incubated with H₂¹⁸O in BHI broth under the same conditions yielded recovered HOIQ which contained no significant amount of ¹⁸O.

ACKNOWLEDGEMENT

This work was supported by grant number CA 40821 from the National Cancer Institute, National Institutes of Health.

REFERENCES

1. (a) T. Sugimura and S. Sato, Cancer Res. (Suppl.), **1983**, 43, 2415s. (b) J. S. Felton, F. T. Hatch, M. G. Knize, and L. F. Bjeldanes, In Diet and Cancer: From Basic Research to Policy Implications; D. A. Roe, Ed.; Alan R. Liss: New York 1983 p. 177. (c) T. Sugimura, Mutation Res., **1985**, 150, 33. (d) H. Kasai, Z. Yamaizumi, S. Nishimura, K. Wakabayashi, M. Nagao, T. Sugimura, N. E. Springarn, J. H. Weisburger, S. Yokoyama, and T. Miyazawa, J. Chem. Soc., Perkin Trans. I, **1981**, 2290. (e) W. A. Hargraves and M. W. Pariza, Cancer Res., **1983**, 43, 1467.
2. (a) E. C. Miller and J. A. Miller, Cancer, **1986** 58, 1795. (b) L. L. Vuolo and G. U. Schuessler, Environ Mutagenesis, **1985**, 7 577. (c) A. J. Miller, Food Technology, **1985**, 39, 75, 109. (d) T. Sugimura and M. Nagao, CRC Critical Reviews in Toxicology, **1979**, 6, 189. (e) T. Sugimura, Cancer, **1982**, 49, 1970.
3. H. Ohgaki, H. Hasegawa, T. Kato, M. Suenaga, M. Ubakata, S. Sato, S. Takayama and T. Sugimura, Proc. Jpn. Acad., **1985**, 61B, 137.
4. S. Takayama, Y. Nakatsuru, M. Masuda, H. Ohgaki, S. Sato and T. Sugimura, Gann, **1984**, 75, 467.
5. W. S. Barnes and J. H. Weisburger, Mutation Res., **1985**, 156, 83.
6. Y. Yamazoe, M. Shimada, T. Kamataki and R. Kato, Cancer Res. **1983**, 43, 5768.
7. M. Bashir, D. G. I. Kingston, R. J. Garman and R. L. van Tassell, T. D. Wilkins, Mutation Res., **1987**, 90, 187.
8. R. J. Carman, R. L. van Tassell, T. D. Wilkins, M. Bashir and D. G. I. Kingston, unpublished results.
9. A. L. Waterhouse and H. Rapoport, J. Labelled Comp., **1985**, 22, 201.

10. E. Ochiai and T. Okamoto, Yakugaku Zasshi, 1948, 68, 88.
11. (a) E. Ochiai and K. Satake, Yakugaku Zasshi, 1951, 71, 1078. (b) Y. Kawazoe and M. Tachibana, Chem. Pharm. Bull., 1967, 15, 1.
12. O. Hayaishi, Science, 1969, 164, 389.
13. (a) R. Hirschberg and J. C. Ensign, J. Bacteriol. 1971, 108 757-759. (b) I. Harary, J. Biol. Chem., 1957, 227 823.
14. W. C. Still, M. Kahn and A. Mitra, J. Org. Chem., 1978, 43, 2923.
15. L. Adolfsson and K. Olsson, Acta Chem. Scand., 1983, 157.
16. R. J. Carman, R. L. Van Tassell, T. D. Wilkins, M. Bashir and D.G.I. Kingston, Abstr. Am. Soc. Microbial. Annual Meeting, 87, 1987, 0-16.

Received, 26th May, 1987