

THE 7-N-OXIDES OF PURINES RELATED TO NUCLEIC ACIDS: THEIR
CHEMISTRY, SYNTHESIS, AND BIOLOGICAL EVALUATION†

Tozo Fujii,*‡ Taisuke Itaya,‡ and Kazuo Ogawa§

‡*Faculty of Pharmaceutical Sciences, Kanazawa University, Takara-machi,
Kanazawa 920, Japan; §Research Laboratories, Ikeda Mohando Co., Ltd.,
Kamiichi-machi, Toyama 930-03, Japan*

Abstract — Recent advances in the chemistry, synthesis, and biological evaluation of the 7-*N*-oxides of purines related to nucleic acids are reviewed. The 7-*N*-oxides covered are those of guanine (1), adenine (2), and hypoxanthine (3) and of related compounds such as 6-mercaptopurine (6-MP) (72), the 6-thioxo analogue of 3, and 6-methylthiopurine, a simple model for azathioprine (78), which were all unknown until recently.

CONTENTS


- I. Introduction
- II. Occurrence
- III. Chemistry and Synthesis
 - A. Guanine 7-Oxide
 - B. Hypoxanthine 7-*N*-Oxide
 - C. Adenine 7-Oxide
- IV. Related Compounds
 - A. 6-Mercaptopurine 7-*N*-Oxide
 - B. 6-Methylthiopurine 7-*N*-Oxide
- V. Biological Activity
- References and Notes

†Dedicated to Emeritus Professor Dr. Shigeru Oae (University of Tsukuba) on the occasion of his 77th birthday.

I. Introduction

Guanine (1) and adenine (2) are important fundamental biomolecules related to DNA's and RNA's. Hypoxanthine (3) is also a biologically significant oxopurine, which occurs in the animal body during the breakdown of nucleic acids and in the plant kingdom as well.¹ It also occurs as the 9- β -D-ribofuranoside inosine and the related nucleotide inosine 5'-phosphate, the former having been identified² as a minor component of more than 30 species of tRNA and the latter being an important precursor in the *de novo* biosynthesis of purine nucleotides such as adenosine 5'-phosphate and guanosine 5'-phosphate.^{1b,3}

Because these three purines carry four endocyclic nitrogen atoms, four kinds of mono-*N*-oxide should be theoretically possible for each. Among the four possible isomeric *N*-oxides⁴ in each case, the 1-, 3-, and 9-*N*-oxides have been prepared by chemical synthesis: 1-hydroxyguanine,⁵ guanine 3-*N*-oxide,^{6,7} 9-hydroxyguanine,⁸ adenine 1-oxide (41),⁹ adenine 3-oxide,¹⁰ 9-hydroxyadenine,¹¹ 1-hydroxyhypoxanthine,¹² hypoxanthine 3-oxide,^{10,13} and 9-hydroxyhypoxanthine.⁸ The remaining 7-*N*-oxide isomers became known only recently, and the definite advances in the chemistry, synthesis, and biological evaluation of the 7-*N*-oxides of these purines emphasize the need for the present review, which covers the literature through the end of 1995.




II. Occurrence


In 1985, three research groups¹⁴⁻¹⁶ independently reported the isolation of guanine 7-oxide (4) from the culture broths of certain *Streptomyces* species (ATCC 39364;¹⁴ *S. purpurascens* A-347;¹⁵ and No. 3780¹⁶), together with its observed antitumor,¹⁴⁻¹⁷ antimicrobial,¹⁶ and antiviral¹⁸ activities. The chemical structure of this antibiotic was established as 4 on the basis of elemental^{14,15b,16} and spectral^{15b,16} analyses; its chemical behavior;¹⁶ and the X-ray molecular structures of the hydrobromide salt (monohydrate),¹⁴ the free base (dihydrate),^{15b} and the pentamethylated derivative (17).¹⁶ Thus, 4 has so far been a unique purine *N*-oxide shown to occur in nature.

III. Chemistry and Synthesis

A. GUANINE 7-OXIDE





Scheme 1








a: R = Me

b: R = MeCH₂CH₂

c: R = CH₂=CH-CH₂



f: R = HO(CH₂)₄



l: R = H

Scheme 2


Direct oxidation of guanine (**1**) with peroxytrifluoroacetic acid has been shown to produce the 3-*N*-oxide,⁶ not the 7-*N*-oxide (**4**) as once thought.¹⁹ In a preliminary experiment, Brown's group²⁰ obtained 8-hydroxyguanine (**6**) from the reaction of 2,4-diamino-5-nitroso-6-hydroxypyrimidine (**5**) with formalin and merely mentioned that it may be possible to obtain the presumed intermediate, guanine 7-*N*-oxide (**4**) (Scheme 1).

The first chemical synthesis of **4** was accomplished by Fujii and co-workers²¹ via a newly devised "phenacylamine route", as delineated in Scheme 2. The route started from coupling of 2-amino-6-chloro-5-nitro-4(3*H*)-pyrimidinone (**7**) with appropriate *N*-substituted phenacylamines, generated *in situ* from the corresponding hydrochlorides (**8a-l**) and 1 N aqueous NaOH, giving the 6-phenacylamino-4-pyrimidinones (**9a-l**). On treatment with 2 N aqueous NaOH at room temperature for 10–60 min, the nitro-pyrimidinones (**9a-k**) cyclized via **10a-k** to provide the 9-substituted guanine 7-oxides (**12a-k**), with elimination of benzoic acid. A similar alkali-treatment of **9l** failed to yield guanine 7-oxide (**4**). However, removal of the 9-(arylmethyl) group from **12i-k** was effected with conc. H₂SO₄ at room temperature for 1–3 h in the presence of toluene, producing the target *N*-oxide (**4**). Application of the same procedure to the unmodified benzyl analogue (**12g**) or the allyl analogue (**12c**) failed to give the desired product (**4**).

As regards the problem of the tautomeric forms of guanine 7-*N*-oxide in the solid state, the N(7)-oxide form (**4**) has been preferred by Kern *et al.*¹⁴ on the basis of the X-ray crystal structure of the corresponding hydrobromide salt monohydrate. On the other hand, Kitahara *et al.*^{15b} have proposed the N(7)-OH form (**11**) on the basis of the result of an X-ray analysis of a single crystal of the dihydrate of the free base grown in 15% tetrahydrofuran–2 M NH₄OH. In solution, the two forms may coexist at equilibrium,¹⁶ and a uv spectroscopic approach, together with three pK_a values (2.6, 5.8, and 9.5) reported¹⁶ for the free base, may suggest that the neutral species of guanine 7-*N*-oxide has a considerable proportion of the N(7)-oxide structure in H₂O (Scheme 2).^{21b}

Scheme 3 summarizes the chemical behavior of **4**. On reduction with Raney Ni, **4** produced guanine (**1**) quantitatively.^{15b} Treatment of **4** with refluxing AcOH gave 8-hydroxyguanine (**6**).¹⁶ The formation of **6** from **4** may be explained by assuming **13** and **14** as the intermediates.²² Permethylation of **4** with MeI in dimethyl sulfoxide (DMSO) in the presence of NaH yielded the pentamethyl derivative (**17**) as the main product.¹⁶ Treatment of **4** with MeI in *N,N*-dimethylacetamide (DMAc) at 25–40°C for 72 h gave 9-methyl-8-hydroxyguanine (**15**) (29% yield), which was identical with a sample prepared from **12a** in 80% yield by treatment with hot AcOH.²² On the other hand, treatment of **4** with dimethyl sulfate in 0.1 N aqueous NaOH furnished 8-methoxyguanine (**16**) (60% yield), which was identical with a sample obtained in 94% yield from 8-methoxyguanosine by glycosidic hydrolysis with *p*-TsOH/AcOH (60–65°C, 40 min).²² Methylation of **6** with dimethyl sulfate in 0.1 N aqueous NaOH failed to give **16**.²² Kitahara *et*

*al.*²³ also observed similar replacement reactions at the 8-position during their attempted chemical modifications of **4**. Werbovetz and Macdonald²⁴ reported that methylation of 9-benzylguanine 7-oxide (**12g**) with trimethyloxonium tetrafluoroborate in MeNO₂ furnished 9-benzyl-7-methoxyguanine salt (**18**) and that **18** was converted to a mixture of 9-benzyl-8-hydroxyguanine (**19**) and 9-benzyl-8-methoxyguanine (**20**) upon treatment with refluxing aqueous NaOH (Scheme 4).²⁵



Several derivatives of **4** have been prepared for biological evaluation. Fujii and co-workers²⁶ synthesized 8-methylguanine 7-oxide (**21**), a model for C(8)-blocked derivatives of **4**, via an α -methylphenacylamine version of the "phenacylamine route" (Scheme 2), which started from condensation of **7** with α -(4-methoxybenzylamino)propiophenone (**22**) and proceeded through cyclization of the resulting phenacylamino-pyrimidinone and removal of the 4-methoxybenzyl group. Nishii *et al.*²⁷ prepared guanosine 7-oxide (**23**) from **4** and ribose 1-phosphate by utilizing purine nucleoside phosphorylase from *Bacillus subtilis* PCI 219. Kitahara *et al.*²⁸ prepared **23** or the 2'-deoxy analogue (**24**) from **4** and ribose 1-phosphate or deoxyribose 1-phosphate using purine nucleoside phosphorylase from bovine spleen. The same group²³ also reported enzymatic conversion of **4** into guanosine 7-oxide 5'-monophosphate disodium salt (**25**) and chemical conversion of **4** into the N²-tetrahydropyranyl derivative (**26**).

B. HYPOXANTHINE 7-N-OXIDE

Hypoxanthine 7-*N*-oxide (**30**) was not known until 1988, when Fujii's group²⁹ achieved its chemical synthesis by extending the "phenacylamine route" (Section III, A) to cover the synthesis of this new purine 7-*N*-oxide at the hypoxanthine level. The first step for the synthesis of **30** was coupling of *N*-(4-methoxybenzyl)phenacylamine, generated *in situ* from the corresponding hydrochloride (**8i**) and 1 N aqueous NaOH, with 6-chloro-5-nitro-4(3*H*)-pyrimidinone (**27**), which was effected in EtOH at room temperature for 6 h to furnish the phenacylamino-pyrimidinone (**28**) (Scheme 5). On treatment with 2 N aqueous NaOH at room temperature for 1 h, **28** gave the *N*-oxide (**31**) and benzoic acid as well. Removal of the 4-


methoxybenzyl group was then carried out with 90% aqueous H_2SO_4 at 30°C for 1 h in the presence of toluene, affording the target 7-*N*-oxide (**30**). Three pK_a values of <1.4 (basic) (for protonated form \rightleftharpoons neutral form), 5.02 (acidic) (for neutral form \rightleftharpoons monoanion), and 10.23 (acidic) (for monoanion \rightleftharpoons dianion) have been obtained spectrophotometrically for **30**,^{29b} and a uv spectroscopic approach has suggested that the neutral form of **30** exists in H_2O mainly as the N(7)-OH tautomer (**29**).^{29b}

The chemical properties of **30** are illustrated in Scheme 6.²⁹ On hydrogenolysis using Raney Ni catalyst and H_2 in H_2O , **30** produced hypoxanthine (**3**). Treatment of **30** with hot AcOH for 20 h or with boiling 2 N aqueous HCl for 1 h gave 6,8-dioxopurine (**32**). The apparent migration of the oxygen function from N(7) to C(8) in this case is analogous to that observed for guanine 7-oxide (**4**) (Section III, A). Methylation of **30** with dimethyl sulfate in 0.2 N aqueous NaOH at room temperature afforded 7-methoxyhypoxanthine (**35**) (28% yield), 7-methoxy-1-methylhypoxanthine (**36**) (7%), and 7-methoxy-3-methylhypoxanthine (**37**) (3%). The locations of the methyl groups were established by reductive demethylations of **35**, **36**, and **37** (Raney Ni/ H_2), which led to the formation of hypoxanthine (**3**), 1-methylhypoxanthine (**33**), and 3-methylhypoxanthine (**34**), respectively. On the other hand, methylation of **30** with MeI in DMAc at room temperature in the absence of alkali gave a complex mixture of products presumed to contain 7-methoxy-9-methyl derivative (**38**), and the mixture yielded 9-methylhypoxanthine (**39**) when subjected to hydrogenolysis (Raney Ni/ H_2) after removal of iodide ion by the use of Dowex 50W-X8 (H^+). The compound (**39**) was identical with a sample prepared from 9-methyladenine (**40**) by deamination with NaNO_2 in aqueous HCl at 90°C .

C. ADENINE 7-OXIDE

Adenine (**2**) has a bicyclic ring system consisting of a 4-aminopyrimidine and an imidazole ring in juxtaposition.³⁰ On treatment with 30% aqueous H_2O_2 in AcOH at room temperature, it undergoes *N*-oxidation preferentially at the 1-position to produce adenine 1-oxide (**41**) in good yield (Scheme 7).^{9,31} This regioselectivity appears to reflect the generalization³² that on *N*-oxidation pyrimidine compounds form only mono-*N*-oxides, whereas imidazoles are resistant to *N*-oxidation.

In 1968, however, Rhaese³³ claimed that treatment of **2** with 0.1 M H_2O_2 in 0.01 M phosphate buffer (pH 7.0) at 37°C for 5 days afforded adenine 7-oxide (**45**) (isolated as a monohydrate sensitive to uv light) in 5% yield without any detectable formation of the N(1)-oxide (**41**). He further claimed that the N(7)-oxide (**45**) was among the products of X-ray irradiation of **2** in 0.05 M phosphate buffer (pH 7.0).³³ Later on, these results were reportedly reproduced by Yamamoto,³⁴ who further asserted that **45** bound noncovalently to urease, an SH protein, in an experiment using a sample of **45** prepared by the method of Rhaese. This unusual regioselectivity of *N*-oxidation of **2** was so striking as to appear questionable. Moreover, the





Scheme 7

chemical and spectroscopic evidence adduced by both authors appeared insufficient to allow definite assignment of the N(7)-oxide structure to their samples, which they thought to be the new *N*-oxide (**45**).


Fujii and co-workers³⁵ reexamined the H₂O₂/buffer oxidation procedure³³ of Rhaese for **2**, but completely failed to reproduce his results; they were unable to obtain any *N*-oxide from **2**. This led them to design a three-step route for the synthesis of adenine 7-oxide (**45**) from adenine (**2**) (Scheme 7).³⁵ Treatment of 3-benzyladenine (**43**), easily obtainable from **2** according to the literature procedure,³⁶ with magnesium monoperoxyphthalate hexahydrate (MMPP·6H₂O) in MeOH at 30°C for 20 h or with *m*-chloroperoxybenzoic acid (MCPBA) in MeOH–1 M acetate buffer (pH 5.0) (1 : 1, v/v) at 30°C for 15 h gave 3-benzyladenine 7-oxide (**44**) in 40% or 24% yield, respectively. The use of 30% aqueous H₂O₂ in AcOH at room temperature or MCPBA in AcOH at 30°C as the oxidizing agent was found to be ineffective. On treatment with conc. H₂SO₄ at 35°C in the presence of toluene for 3 h, **44** furnished the desired compound, adenine 7-oxide (**45**), in 55% yield. Characterization of **45** as the N(7)-oxide was readily achieved by measurement of its uv spectrum, which was different from those of the three known isomeric *N*-oxides, and by its chemical reactions including deamination and methylation, as shown in Scheme 7. In addition, the location of the oxygen function in **44** and **45** was confirmed by X-ray crystallographic analysis.^{35b}

Fujii's group^{35b} further found that treatment of **43** with a large excess of 30% aqueous H₂O₂ in MeOH in the presence of MeCN and KHCO₃ at 25°C for 22 h produced the N(7)-oxide (**44**) and 7-acetamido-3-benzyladenine (**50**) in 12% and 1% yields, respectively, together with 28% recovery of **43**. They determined p*K*_a values of **45** spectrophotometrically in H₂O at 30°C, obtaining two values of 3.4 (basic) [for protonated form (**51**) ⇌ neutral form] and 5.75 (acidic) [for neutral form ⇌ monoanion (**53**)] (Scheme 8).^{35b} A uv spectroscopic approach suggested that the neutral species of **45** exists in H₂O as an equilibrated mixture of the N(7)-oxide (**45**) and N(7)-OH (**52**) tautomers.^{35b} As in the case of 3-benzyladenine (**43**) described above, 3-methyladenine (**48**) and 3-ethyladenine (**54**) underwent peroxycarboxylic acid oxidation at N(7), giving **47** and **55** in 13–25% yields.^{37a} Treatment of **44**, **47**, and **55** with alkyl halide (R²X) in DMAc at 30°C afforded the corresponding 7-alkoxy derivatives (**56**) (81–91% yields), which furnished 3-alkyl-8-hydroxyadenines (**57**) in 26–50% yields on treatment with 0.1 N aqueous NaOH at room temperature (Scheme 9).^{37a} Treatment of **56** (X = ClO₄) with 1 M R³ONa in R³OH (R³ = Me, Et, or PhCH₂) at room temperature gave 8-alkoxy-3-alkyladenines (**58**) in 28–97% yields, and hydrolysis of **58** (R³ = Me) with boiling 1 N aqueous HCl or hydrogenolysis (Pd–C/H₂) of **58** (R³ = PhCH₂) provided **57** in 73–88% yields.^{37b}


Much information has been accumulated concerning the regioselectivity in *N*-oxidation of *N*^x-benzyladenines. Oxidation of 9-benzyladenine (**59**) with peroxyacetic acid gives the N(1)-oxide (**63**) in 69% yield;³⁸ that of 9-benzyladenine-2-*d* (**60**) with MCPBA gives the corresponding N(1)-oxide (**64**) in 71%



Scheme 8



Scheme 9



Scheme 10

yield;³¹ oxidation of 7-benzyladenine (**61**) with MCPBA affords the N(1)-oxide (**65**) in 76% yield;³⁹ oxidation of 3-benzyladenine (**43**) with MMPP or with MCPBA or with aqueous H₂O₂/MeCN/KHCO₃ provides the N(7)-oxide (**44**) in 40% or 24% or 12% yield, respectively (*vide supra*);³⁵ and oxidation of N⁶-benzyladenine (**62**) with MCPBA⁴⁰ or with trifluoroperoxyacetic acid⁴¹ furnishes the N(1)-oxide (**66**) (35% yield) or the N(3)-oxide (4%) and N(7)-oxide (4%), respectively. Fujii's group⁴² treated 1-benzyladenine (**67**), the remaining positional isomer, with MCPBA in MeOH or in MeOH–0.5 M phosphate buffer (pH 6.6) at 30°C and obtained 1-benzyladenine 7-oxide (**68**) as the main product (Scheme 10). Nonreductive debenylation of **68** with H₂SO₄/toluene gave adenine 7-oxide (**45**) in 63% yield. The structure of **68** was unequivocally established by an X-ray crystallographic analysis. Thus, the reaction sequence **67** → **68** → **45** afforded an alternative synthesis of **45**.⁴²


Yet another synthetic approach to **45** would be an extension of the "phenacylamine route" (Section III, A), as shown in Scheme 10. Fujii's group⁴² obtained **71** in 67% yield from 4-amino-6-chloro-5-nitropyrimidine (**70**) and **8i**. However, treatment of **71** with 2 N aqueous NaOH in MeOH at room temperature for 2 h gave a mixture of many products, from which they were unable to isolate the cyclized product (**69**), even if it were present. This led them to abandon the "phenacylamine route" approach.

IV. Related Compounds

A. 6-MERCAPTOPURINE 7-N-OXIDE

6-Mercaptopurine (6-MP) (**72**), the 6-thio analogue of hypoxanthine (**3**), is an antileukemic agent of longstanding clinical usefulness.⁴³ Among the four possible *N*-oxides of 6-MP, only the N(3)-oxide (**75**) has so far been obtained. It has been synthesized from 6-chloropurine 3-oxide (**76**) and ammonium dithiocarbamate⁴⁴ or from 7-aminothiazolo[5,4-*d*]pyrimidine 6-*N*-oxide (**77**) by rearrangement,⁴⁵ and a comparison of the activities of the *N*-oxide (**75**) with the parent 6-MP has been made in several biological systems.⁴⁵ Fujii *et al.*⁴⁶ reported the first synthesis of 6-mercaptopurine 7-*N*-oxide (**73**), in which they adopted a dichloropyrimidine variant (Scheme 11) of their favorite "phenacylamine route" (Section III, A and B). The synthesis of **73** started with the condensation of *N*-(4-methoxybenzyl)phenacylamine, generated from its hydrochloride salt (**8i**), with 4,6-dichloro-5-nitropyrimidine (**83**) in CHCl₃ at 0–5°C for 1 h to give the phenacylaminopyrimidine (**79**). Successive treatments of **79** with thiourea, conc. aqueous NH₃, and 2 N aqueous NaOH afforded the *N*-oxide (**80**). Removal of the 4-methoxybenzyl group from **80** was effected in a mixture of conc. H₂SO₄ and toluene at 23°C for 2 h, giving the target compound (**73**). On treatment with sodium dithionite in boiling 50% aqueous MeOH for 1.5 h, **73** produced 6-MP (**72**). The ¹H nmr spectrum of **73** indicated that 6-mercaptopurine 7-*N*-oxide exists in Me₂SO-*d*₆ in the 6-thio-1*H*-


purine form (**85**) rather than the C(6)-SH form (**73**). In an alternative synthetic approach to **73**, Fujii *et al.*⁴⁶ heated hypoxanthine 7-*N*-oxide (**30**) with P₂S₅ in boiling pyridine. However, they were unable to obtain **73**, but isolated a compound inferred to be 8-mercaptopyoxanthine. They obtained three p*K*_a values for **73** spectrophotometrically in H₂O at 30°C, as shown in Scheme 12, and a uv spectroscopic approach suggested the overwhelming predominance of the N(7)-OH tautomer (**88**) over the N(7)-oxide tautomer (**73**) in the neutral species of 6-mercaptapurine 7-*N*-oxide in H₂O.^{46b}



Scheme 12

B. 6-METHYLTHIOPURINE 7-*N*-OXIDE

Azathioprine (Imuran[®]) (**78**), the *S*-(1-methyl-4-nitro-1*H*-imidazol-5-yl) derivative of 6-MP (**72**), is an immunosuppressive agent of longstanding clinical usefulness.⁴³ It acts as a pro-drug for 6-MP.^{43a} Fujii *et al.*⁴⁶ reported the first synthesis of 6-methylthiopurine 7-*N*-oxide (**74**), a simple model for the 7-*N*-oxide of azathioprine (**78**). For the synthesis of **74**, they methylated the precursor (**80**) for **73** with dimethyl sulfate in a mixture of 1 N aqueous NaOH and MeOH at room temperature or with MeI and K₂CO₃ in MeOH at room temperature, obtaining the 6-methylthio derivative (**81**) (Scheme 11). Nonreductive debenzoylation of **81** with conc. H₂SO₄ in the presence of toluene at 25°C for 1 h gave the desired *N*-oxide (**74**). On the other hand, direct methylation of **73** with MeI or dimethyl sulfate in a mixture of MeOH and 1 N aqueous NaOH resulted in the formation of a mixture of many products, from which they were unable to obtain the *S*-methyl derivative (**74**). The location of the oxygen function in **73**, **74**, and **80** was confirmed by X-ray crystallographic analysis of **74**·H₂O, which was shown to exist in the N(7)-OH form (**86**).



Scheme 13

Fujii *et al.*^{46b} also determined two pK_a values for **74** spectrophotometrically in H₂O at 30°C, as shown in Scheme 13, and a uv spectroscopic approach suggested that the neutral species of 6-methylthiopurine 7-*N*-oxide exists in H₂O as an equilibrated mixture of the N(7)-oxide (**74**) and the N(7)-OH (**86**) tautomers. In an attempt to develop an alternative synthetic route to adenine 7-oxide (**45**), Fujii *et al.*⁴⁶ examined amination of **74** under a variety of reaction conditions (Scheme 11). However, all attempts resulted in the recovery of **74**, suggesting the inertness of the C(6)-SMe group in the anionic species (**92**). On the other hand, treatment of the N(9)-arylmethyl derivative (**81**) with 16% methanolic NH₃ at 24°C for 4 h gave an unstable crude compound inferred to be the ring-opened product (**82**), which reverted to **81** on heating in boiling EtOH for 30 min.⁴⁶ Treatment of **81** with saturated ethanolic NH₃ in an autoclave at 110°C for 6 h afforded the C(8)-amino derivative (**84**).⁴⁶ In either case, the desired adenine derivative (**69**) could not be obtained.

V. Biological Activity

Guanine 7-oxide (**4**) exhibited excellent activity in mice that were inoculated either intraperitoneally or subcutaneously with L1210 leukemia cells.^{14,15a} It caused pronounced growth inhibition of several murine and human cell lines *in vitro*.¹⁷ For the L1210 lymphoblastic leukemia, 50% inhibition was obtained below 1 μ M.¹⁷ It also inhibited Yoshida sarcoma and L5178Y leukemia cells in culture at IC₅₀'s of 1.65 and 2.40 μ g/ml, respectively.¹⁶ The intraperitoneal administration of this antibiotic showed a life-prolongation effect on mice bearing P388 leukemia, and the activity at a dose of 6.0 mg/kg/day was almost comparable to that observed with mitomycin C at a dose of 1.0 mg/kg/day.¹⁶ The *N*-oxide (**4**) also showed a dose-dependent inhibition of the growth of Ehrlich solid carcinoma in mice *via* oral administration.¹⁶ Guanine 7-oxide (**4**) has no activity against *Staphylococcus aureus* 209P, *Bacillus subtilis* PCI 219, *Escherichia coli* NIHJ, *Pseudomonas aeruginosa* IFO 3445, *Micrococcus luteus* PCI 1001, *Candida albicans* 3147, and *Saccharomyces cerevisiae* at a concentration of 100 μ g/ml.^{15a} Nishii *et al.*¹⁶ reported that **4** was inhibitory to *Candida albicans* but inactive against Gram-positive and Gram-negative bacteria and *Trichophyton* species. Moderate antiviral activity of **4** was demonstrated against DNA and RNA viruses derived from salmonids.¹⁸ The LD₅₀ of **4** in mice was determined to be 40–80 mg/kg (by single intraperitoneal administration)^{15a} or 53 mg/kg (by intraperitoneal administration).¹⁶ Jackson *et al.*¹⁷ reported that the *N*-oxide (**4**) is converted within sensitive cells into guanosine 7-oxide 5'-triphosphate and this results in inhibition of cellular protein synthesis. Nishii *et al.*²⁷ reported that the antimicrobial activity of guanosine 7-oxide (**23**) was very weak but it inhibited L5178Y mouse leukemia cells in culture at an IC₅₀ of 0.60 μ g/ml; intraperitoneal administration of **23** showed a life-prolongation effect on mice bearing P388 leukemia; and **23** showed a dose-dependent inhibition of the growth of Ehrlich solid carcinoma in mice. Kitahara *et al.*²⁸

reported the antitumor activities of **23** and the 2'-deoxy analogue (**24**) against mouse leukemia L1210 cells. They also reported the biological activities of the nucleotide analogue (**25**) and the *N*²-tetrahydropyranyl derivative (**26**).²³

In the *in vitro* bioassay of antileukemic activity against murine L5178Y cells, Fujii and co-workers^{21b} found that none of the 9-substituted guanine 7-oxide (**12a-k**) was more effective than the parent, natural *N*-oxide (**4**). Within this series, however, the benzyl analogues (**24g-k**) with or without alkoxy functions were more cytotoxic, with IC₅₀'s of 13.0–48.0 µg/ml, than the alkyl analogues (**12a-f**). 8-Methylguanine 7-oxide (**21**), 9-(4-methoxybenzyl)-8-methylguanine 7-oxide, and 9-(4-methoxy-3-sulfobenzyl)-8-methylguanine 7-oxide showed only weak antileukemic activity and no antimicrobial activity.²⁶

Hypoxanthine 7-*N*-oxide (**30**) was weakly cytotoxic, with IC₅₀ of 100 µg/ml, in the *in vitro* bioassay of antileukemic activity against murine L5178Y cells, and it did not show any antimicrobial activity even at 1000 µg/ml.^{29b} None of its 9-(4-methoxybenzyl) derivative (**31**) and the 7-methoxy derivatives (**35**, **36**, and **37**) was found to be antileukemic or antimicrobial.^{29b} In a similar bioassay, 6-mercaptopurine 7-*N*-oxide (**73**) and its 9-(4-methoxybenzyl) derivative (**80**) were less effective than the parent 6-MP (**72**), but slightly more cytotoxic than hypoxanthine 7-*N*-oxide (**30**); 6-methylthiopurine 7-*N*-oxide (**74**) and adenine 7-oxide (**45**) were inactive at 50 µg/ml concentration.^{46b}

In the tobacco callus bioassay of cytokinin activity, each of *N*⁶-benzyladenine 1-oxide (**66**), *N*⁶-benzyladenine 3-oxide, and *N*⁶-benzyladenine 7-oxide was active at 4 µM concentration, being less active than the parent synthetic cytokinin (**62**) by a factor of 40.⁴¹

REFERENCES AND NOTES

1. For pertinent reviews, see (a) J. H. Lister, 'Fused Pyrimidines. Part II: Purines,' ed. by D. J. Brown, Wiley-Interscience, New York, 1971, pp. 250–251; (b) G. Shaw, 'Rodd's Chemistry of Carbon Compounds,' 2nd ed., Vol. IV, Part L, ed. by S. Coffey, Elsevier Scientific Publishing Co., Amsterdam, 1980, pp. 31–32; (c) Atta-ur-Rahman and M. I. Choudhary, 'The Alkaloids,' Vol. 38, ed. by A. Brossi, Academic Press, New York, 1990, pp. 256–257.
2. For a review, see M. Sprinzl, T. Hartmann, J. Weber, J. Blank, and R. Zeidler, *Nucleic Acids Res.*, 1989, **17**, r1.
3. For reviews, see (a) Ref. 1a, pp. 112–113; (b) G. Shaw, 'Comprehensive Heterocyclic Chemistry,' Vol. 5, ed. by K. T. Potts, Pergamon Press, Oxford, 1984, p. 505 and pp. 598–599.
4. The designation "*N*-oxide" used in this review follows that adopted in ref. 7, footnote 15.
5. A. A. Watson, S. C. Nesnow, and G. B. Brown, *J. Org. Chem.*, 1973, **38**, 3046.
6. U. Wölcke and G. B. Brown, *J. Org. Chem.*, 1969, **34**, 978.

7. J. C. Parham, T. G. Winn, and G. B. Brown, *J. Org. Chem.*, 1971, **36**, 2639.
8. A. A. Watson, *J. Org. Chem.*, 1974, **39**, 2911.
9. (a) M. A. Stevens, D. I. Magrath, H. W. Smith, and G. B. Brown, *J. Am. Chem. Soc.*, 1958, **80**, 2755; (b) M. A. Stevens and G. B. Brown, *ibid.*, 1958, **80**, 2759; (c) T. Fujii and T. Itaya, *Tetrahedron*, 1971, **27**, 351.
10. I. Scheinfeld, J. C. Parham, S. Murphy, and G. B. Brown, *J. Org. Chem.*, 1969, **34**, 2153.
11. A. A. Watson, *J. Org. Chem.*, 1977, **42**, 1610.
12. (a) E. C. Taylor, C. C. Cheng, and O. Vogl, *J. Org. Chem.*, 1959, **24**, 2019; (b) J. C. Parham, J. Fissekis, and G. B. Brown, *ibid.*, 1966, **31**, 966; (c) H. Kawashima, T. Meguro, and I. Kumashiro, *Bull. Chem. Soc. Jpn.*, 1966, **39**, 633.
13. H. Kawashima and I. Kumashiro, *Bull. Chem. Soc. Jpn.*, 1969, **42**, 750.
14. D. L. Kern, G. C. Hokanson, J. C. French, and N. K. Dally, *J. Antibiot.*, 1985, **38**, 572.
15. (a) M. Kitahara, K. Ishii, Y. Kumada, T. Shiraishi, T. Furuta, T. Miwa, H. Kawaharada, and K. Watanabe, *J. Antibiot.*, 1985, **38**, 972; (b) M. Kitahara, K. Ishii, H. Kawaharada, K. Watanabe, T. Suga, T. Hirata, and S. Nakamura, *ibid.*, 1985, **38**, 977.
16. M. Nishii, J. Inagaki, F. Nohara, K. Isono, H. Kusakabe, K. Kobayashi, T. Sakurai, S. Koshimura, S. K. Sethi, and J. A. McCloskey, *J. Antibiot.*, 1985, **38**, 1440.
17. R. C. Jackson, T. J. Boritzki, J. A. Besserer, K. L. Hamelehle, J. L. Shillis, W. R. Leopold, and D. W. Fry, *Adv. Enzyme Regul.*, 1987, **26**, 301.
18. (a) M. Hasobe, M. Saneyoshi, and K. Isono, *J. Antibiot.*, 1985, **38**, 1581; (b) *Idem, ibid.*, 1986, **39**, 1291.
19. T. J. Delia and G. B. Brown, *J. Org. Chem.*, 1966, **31**, 178.
20. G. Zvilichovsky and G. B. Brown, *J. Org. Chem.*, 1972, **37**, 1871.
21. (a) F. Nohara, M. Nishii, K. Ogawa, K. Isono, M. Ubukata, T. Fujii, T. Itaya, and T. Saito, *Tetrahedron Lett.*, 1987, **28**, 1287; (b) K. Ogawa, M. Nishii, J. Inagaki, F. Nohara, T. Saito, T. Itaya, and T. Fujii, *Chem. Pharm. Bull.*, 1992, **40**, 343.
22. K. Ogawa, Ph. D. Dissertation, Kanazawa University, May 1994, pp. 29–31.
23. M. Kitahara, K. Shinjyo, M. Fukae, K. Hosoe, T. Shiraishi, K. Ishii, K. Watanabe, and H. Kawaharada, *J. Antibiot.*, 1990, **43**, 352.
24. K. A. Werbovetz and T. L. Macdonald, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 2323.
25. For a discussion of similar activation of C(8), see W. Adam and A. Treiber, *J. Am. Chem. Soc.*, 1995, **117**, 2686.

26. K. Ogawa, M. Nishii, J. Inagaki, F. Nohara, T. Saito, T. Itaya, and T. Fujii, *Chem. Pharm. Bull.*, 1992, **40**, 1315.
27. M. Nishii, Y. Fujihara, J. Inagaki, K. Ogawa, F. Nohara, M. Ubukata, and K. Isono, *Agric. Biol. Chem.*, 1986, **50**, 2697.
28. M. Kitahara, K. Ishii, T. Okazaki, T. Hidaka, and K. Watanabe, *J. Antibiot.*, 1986, **39**, 1288.
29. (a) K. Ogawa, T. Saito, F. Nohara, M. Nishii, T. Itaya, and T. Fujii, *Heterocycles*, 1988, **27**, 885; (b) K. Ogawa, M. Nishii, F. Nohara, T. Saito, T. Itaya, and T. Fujii, *Chem. Pharm. Bull.*, 1992, **40**, 612.
30. Ref. 1a, Chapter 1.
31. For a similar *N*-oxidation of adenine-2-*d* to form adenine-2-*d* 1-oxide, see T. Fujii, T. Saito, K. Kizu, H. Hayashibara, Y. Kumazawa, S. Nakajima, and T. Fujisawa, *Chem. Pharm. Bull.*, 1991, **39**, 301.
32. E. Ochiai, 'Aromatic Amine Oxides,' Elsevier, Amsterdam, 1967, pp. 50–51.
33. H.-J. Rhaese, *Biochim. Biophys. Acta*, 1968, **166**, 311.
34. O. Yamamoto, *J. Radiat. Res.*, 1980, **21**, 239.
35. (a) T. Fujii, K. Ogawa, T. Saito, K. Kobayashi, and T. Itaya, *Heterocycles*, 1994, **38**, 477; (b) T. Fujii, K. Ogawa, T. Saito, K. Kobayashi, T. Itaya, T. Date, and K. Okamura, *Chem. Pharm. Bull.*, 1995, **43**, 53.
36. T. Fujii, G. C. Walker, N. J. Leonard, D. C. DeLong, and K. Gerzon, *J. Med. Chem.*, 1979, **22**, 125.
37. (a) T. Fujii, Y. Takada, K. Ogawa, and T. Itaya, Abstracts of Papers, Part 2, 115th Annual Meeting of Pharmaceutical Society of Japan, Sendai, March 1995, p. 73; (b) T. Fujii, Y. Takada, and T. Itaya, Abstracts of Papers, Part 2, 116th Annual Meeting of Pharmaceutical Society of Japan, Kanazawa, March 1996, p. 139.
38. T. Fujii, C. C. Wu, and T. Itaya, *Chem. Pharm. Bull.*, 1971, **19**, 1368.
39. (a) K. Ogawa, T. Saito, T. Itaya, and T. Fujii, *Heterocycles*, 1994, **38**, 253; (b) T. Fujii, K. Ogawa, T. Saito, and T. Itaya, *Chem. Pharm. Bull.*, 1995, **43**, 328.
40. (a) T. Fujii, K. Ogawa, and T. Itaya, *Heterocycles*, 1994, **37**, 219; (b) T. Itaya, K. Ogawa, Y. Takada, and T. Fujii, *Chem. Pharm. Bull.*, 1996, **44**, in press.
41. T. Fujii, Y. Takada, K. Ogawa, T. Itaya, and S. Matsubara, *Chem. Pharm. Bull.*, 1995, **43**, 325.
42. T. Fujii, K. Ogawa, T. Saito, T. Itaya, T. Date, and K. Okamura, *Chem. Pharm. Bull.*, 1995, **43**, 321.

43. (a) G. B. Elion, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 870; (b) T. A. Krenitsky, W. W. Hall, J. L. Selph, J. F. Truax, and R. Vinegar, *J. Med. Chem.*, 1989, **32**, 1471.
44. A. Giner-Sorolla, *J. Heterocycl. Chem.*, 1971, **8**, 651.
45. G. B. Brown, G. Levin, S. Murphy, A. Sele, H. C. Reilly, G. S. Tarnowski, F. A. Schmid, M. N. Teller, and C. C. Stock, *J. Med. Chem.*, 1965, **8**, 190.
46. (a) K. Ogawa, T. Itaya, and T. Fujii, *Heterocycles*, 1994, **38**, 1225; (b) T. Fujii, K. Ogawa, T. Itaya, T. Date, J. Inagaki, and F. Nohara, *Chem. Pharm. Bull.*, 1995, **43**, 408.

Received, 19th February, 1996