

HETEROCYCLES, Vol. 78, No. 4, 2009, pp. 977 - 982. © The Japan Institute of Heterocyclic Chemistry  
Received, 20th October, 2008, Accepted, 27th November, 2008, Published online, 3rd December, 2008  
DOI: 10.3987/COM-08-11584

## **SYNTHESIS OF 4-SUBSTITUTED 3,5-DICYANO-2,6-PIPERIDINE-DIONES USING LITHIUM NITRIDE AS A CONVENIENT SOURCE OF AMMONIA**

**Liqiang Wu, \* Chunguang Yang, Liming Yang, and Lijuan Yang**

School of Pharmacy, Xinxiang Medical College, Xinxiang, Henan 453003, P. R. China. E-mail: wliq1974@sohu.com

**Abstract** – A simple and efficient one-pot synthesis of 4-substituted-3,5-dicyano-2,6-piperidinediones was achieved in good yields via the three-component reaction of aldehyde or ketone, ethyl cyanoacetate, lithium nitride ( $\text{Li}_3\text{N}$ ) as a convenient source of ammonia in MeOH.

### **INTRODUCTION**


4-Substituted-3,5-dicyano-2,6-piperidinediones exhibit a large range of biological activities such as anticonvulsant, sedative and analgesic activities.<sup>1</sup> They are also useful synthetic intermediates for various pharmaceuticals and active compounds.<sup>2</sup> There are two strategies for synthesis of 4-substituted-3,5-dicyano-2,6-piperidinediones: (1) They are synthesized by Guareschi-Thorpe reaction in following manner: The one-pot, three-component condensation of an aldehyde or ketone with ethyl cyanoacetate and ammonia in alcohol.<sup>2-3</sup> The reaction requires at least 48 hours to proceed to completion, and more typically between 48-168 hours for completion. In addition, current environmental regulations make it impractical to produce gaseous  $\text{NH}_3$  in ethanol on a large scale. (2) The other approaches for the preparation of 4-substituted-3,5-dicyano-2,6-piperidinediones using aldehyde or ketone, substituted  $\alpha$ -cyanoacetamide, ethyl cyanoacetate in an ethanolic solution of sodium ethoxide at room temperature have also been reported.<sup>1</sup> Substituted  $\alpha$ -cyanoacetamide as the starting material of the route is not easy to obtain. We now report a simple and efficient route to 4-substituted-3,5-dicyano-2,6-piperidinediones using  $\text{Li}_3\text{N}$  as a convenient source of ammonia.

### **RESULTS AND DISCUSSION**

A range of 4-substituted-3,5-dicyano-2,6-piperidinediones was synthesized from a combination of aldehyde or ketone (**1**), ethyl cyanoacetate (**2**) and  $\text{Li}_3\text{N}$  (**3**) in a 1: 2: 2 ratio in an MeOH solution (Table 1). The reaction was completed 12-16 hours at room temperature and the crude product was isolated by precipitation upon addition of hydrochloric acid to the mixture, the separated solid was filtered off and

crystallized from the appropriate solvent.

**Table 1** Synthesis of 4-substituted-3,5-dicyano-2,6-piperidinediones<sup>a</sup>




Entry	Aldehyde or Ketone	Product 4	Time (h)	Yield (%) <sup>b</sup>
4a			12	79
4b			12	81
4c			16	76
4d			14	82
4e			14	83
4f			14	76
4g			12	84
4h			12	74
4i			12	78
4j			16	75

<sup>a</sup>Aldehyde or ketone: ethyl cyanoacetate: Li<sub>3</sub>N = 1: 2: 2; reactions executed in a sealed vessel at room temperature.

<sup>b</sup>Isolated yield

In recent study, we have discovered that  $\text{Li}_3\text{N}$  as electrode material<sup>4</sup> and catalyst in synthesis of *c*BN<sup>5</sup> in chemistry industry can release ammonia upon treatment with MeOH along with the corresponding lithium methoxide. Thus, this reagent can serve a dual role as it releases ammonia in situ while generating lithium methoxide with the potential to act as a catalyst (see Figure 1).



**Figure 1.** The mechanism of reaction using  $\text{Li}_3\text{N}$  as a source of ammonia

In order to determine the efficiency of  $\text{Li}_3\text{N}$  as a source of ammonia, the same reactions were carried out with ammonia,  $\text{NH}_4\text{Ac}$  and other commercially nitrides ( $\text{Mg}_3\text{N}_2$ ,  $\text{Zn}_3\text{N}_2$ ,  $\text{AlN}$ ) employed in related condensations. The results summarized in Table 2 revealed that these systems gave lower yields of the desired product or no desired product. The results also emphasized the importance of using lithium methoxide (entry 3), which could improve yield.

**Table 2** Synthesis of 4,4-dimethyl-3,5-dicyano-2,6-piperidinediones using different source of ammonia<sup>a</sup>

Entry	Source of Ammonia	Solvent	Yield (%) <sup>b</sup>
1	2 equiv $\text{NH}_3$	$\text{H}_2\text{O}$	6
2	2 equiv $\text{NH}_3$	MeOH	24
3	2 equiv $\text{NH}_3$ +6 equiv $\text{LiOMe}$	MeOH	58
4	2 equiv $\text{NH}_4\text{Ac}$	MeOH	0
5	1 equiv $\text{Mg}_3\text{N}_2$	MeOH	31
6	1 equiv $\text{Zn}_3\text{N}_2$	MeOH	0
7	2 equiv $\text{AlN}$	MeOH	0

<sup>a</sup> Acetone: ethyl cyanoacetate=1: 2, reactions executed in a sealed vessel at room temperature for 12 h.

<sup>b</sup> Isolated yield

In summary, we first developed a simple and efficient methodology to synthesise a range of 4-substituted-3,5-dicyano-2,6-piperidinediones using  $\text{Li}_3\text{N}$  as a convenient source of ammonia. We believe that this methodology will be a valuable addition to the existing methods in the field of 4-substituted-3,5-dicyano-2,6-piperidinediones.

## EXPERIMENTAL

NMR spectra were determined on FT-NMR Avance 400 spectrometer, coupling constants ( $J$ ) were measured in Hz; Elemental analysis were recorded on a PEA-1110 elemental analyzer. Melting points were determined on a Mel-Temp capillary tube apparatus and were uncorrected; Commercially available reagents were used throughout without further purification unless otherwise stated.

**General Procedure for the Preparation of 4.** To a stirred solution of aldehyde or ketone (10 mmol) and ethyl cyanoacetate (2.26 g, 20 mmol) in MeOH (5 mL) at 8 °C in a 10 mL, lithium nitride (0.7 g, 20 mmol) was added in a single portion. The reaction mixture was sealed immediately and stirred at rt for 12-16 h. The mixture was then acidified by hydrochloric acid (15mL in 50 mL water) and the separated solid was filtered off and crystallized from MeOH to give **4** as a white solid.

**4,4-Dimethyl-3,5-dicyano-2,6-piperidinedione (4a)** : mp 215-217°C (lit.,<sup>2b</sup> mp 214-216°C); IR (KBr)  $\nu$ : 3211 (NH), 2242 (CN), 1723, 1698 (C=O)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$ : 12.02 (s, 1 H, NH), 4.42 (s, 2 H, CH), 1.28 (t,  $J=7.5$  Hz, 6 H,  $\text{CH}_3$ ); Anal. Calcd for  $\text{C}_9\text{H}_9\text{N}_3\text{O}_2$ : C, 56.54; H, 4.74; N, 21.98. Found: C, 56.49; H, 4.70; N, 21.89.

**4-Ethyl-4-methyl-3,5-dicyano-2,6-piperidinedione (4b)** : mp 188-190°C (lit.,<sup>2b</sup> mp 191-193°C); IR (KBr)  $\nu$ : 3240 (NH), 2225 (CN), 1743, 1702 (C=O)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$ : 11.92 (s, 1 H, NH), 4.51 (s, 2 H, CH), 1.28-1.23 (m, 2H,  $\text{CH}_2$ ), 1.16 (t,  $J=7.3$  Hz, 3H,  $\text{CH}_3$ ), 0.96-0.92 (m, 3H,  $\text{CH}_3$ ); Anal. Calcd for  $\text{C}_{10}\text{H}_{11}\text{N}_3\text{O}_2$ : C, 58.53; H, 5.40; N, 20.48. Found: C, 58.44; H, 5.45; N, 20.39.

**4-Methyl-4-(4-methylpent-3-enyl)-3,5-dicyano-2,6-piperidinedione (4c)** : mp 192-193°C; IR (KBr)  $\nu$ : 3235 (NH), 2240 (CN), 1730, 1698 (C=O)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$ : 12.08 (s, 1 H, NH), 5.12 (m, 1 H, CH=), 4.43 (s, 2 H, CH), 2.10-2.06 (m, 2H,  $\text{CH}_2$ ), 1.29-1.25 (m, 2H,  $\text{CH}_2$ ), 1.20 (t,  $J=7.2$  Hz, 3H,  $\text{CH}_3$ ), 0.92-0.89 (m, 3H,  $\text{CH}_3$ ); Anal. Calcd for  $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}_2$ : C, 64.85; H, 6.61; N, 16.20. Found: C, 64.70; H, 6.75; N, 16.35.

**6, 10-Dicyano-8-azaspiro[4,5]decane-7,9-dione (4d)** : mp 162-163°C (lit.,<sup>1</sup> mp 166 °C); IR (KBr)  $\nu$ : 3212 (NH), 2243 (CN), 1745, 1708 (C=O)  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$ : 11.94 (s, 1 H, NH), 4.63 (s, 2 H, CH), 1.75-1.69(m, 8H,  $\text{CH}_2$ ); Anal. Calcd for  $\text{C}_{11}\text{H}_{11}\text{N}_3\text{O}_2$ : C, 60.82; H, 5.10; N, 19.34. Found: C, 60.84; H, 5.15; N, 19.41.

**1, 5-Dicyano-3-azaspiro[5,5]undecane-2,4-dione (4e)** : mp 196-198°C (lit.,<sup>1</sup> mp 200 °C); IR (KBr)  $\nu$ : 3201(NH), 2239(CN), 1732, 1712 (C=O)  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$ : 12.02 (s, 1 H, NH),

4.53 (s, 2 H, CH), 1.69-1.43(m, 10H, CH<sub>2</sub>); Anal. Calcd for C<sub>12</sub>H<sub>13</sub>N<sub>3</sub>O<sub>2</sub>: C, 62.33; H, 5.67; N, 18.17. Found: C, 62.41; H, 5.60; N, 19.21.

**7,11-Dicyano 3-Oxa-9-azaspiro[5.5]undecane-8,10-dione (4f)** : mp 196-198°C. IR (KBr)  $\nu$ : 3234 (NH), 2219 (CN), 1734, 1705 (C=O) 1185 (C-O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$ : 12.10 (s, 1 H, NH), 4.92 (s, 2 H, CH), 3.72-3.66 (m, 4 H, OCH<sub>2</sub>), 1.72-1.68 (m, 4H, CH<sub>2</sub>); Anal. Calcd for C<sub>11</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>: C, 56.65; H, 4.76; N, 18.02. Found: C, 56.45; H, 4.68; N, 18.15.

**4-Phenyl-3,5-dicyano-2,6-piperidinedione (4g)** : mp 248-249°C (lit.,<sup>1</sup> mp 255°C); IR (KBr)  $\nu$ : 3305 (NH), 2260 (CN), 1712, 1692 (C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$ : 12.02 (s, 1 H, NH), 7.45-7.32 (m, 5 H, ArH), 4.96-4.87 (m, 2 H, CH), 4.40-4.32 (m, 1 H, CH); Anal. Calcd for C<sub>13</sub>H<sub>9</sub>N<sub>3</sub>O<sub>2</sub>: C, 52.79; H, 5.64; N, 22.39. Found: C, 52.65; H, 5.58; N, 22.35.

**4-(4-Fluorophenyl)-3,5-dicyano-2,6-piperidinedione (4h)** : mp 245-247°C (lit.,<sup>1</sup> mp 255°C); IR (KBr)  $\nu$ : 3315 (NH), 2274 (CN), 1726, 1702 (C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$ : 12.08 (s, 1 H, NH), 7.20-6.92(m, 4 H, ArH), 4.73-4.69 (m, 2 H, CH), 4.34-4.28 (m, 1 H, CH); Anal. Calcd for C<sub>13</sub>H<sub>8</sub>FN<sub>3</sub>O<sub>2</sub>: C, 60.70; H, 3.13; N, 16.33. Found: C, 60.45; H, 3.15; N, 16.20.

**4-(4-Methoxyphenyl)-3,5-dicyano-2,6-piperidinedione (4i)** : mp 317-318°C (lit.,<sup>1</sup> mp 325°C); IR (KBr)  $\nu$ : 3294 (NH), 2256 (CN), 1736, 1698 (C=O), 1282, 1088 (C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$ : 12.02 (s, 1 H, NH), 7.40-6.95(m, 4 H, ArH), 4.82-4.76 (m, 2 H, CH), 4.22-4.16 (m, 1 H, CH), 3.74 (s, 3 H, CH<sub>3</sub>); Anal. Calcd for C<sub>14</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>: C, 62.45; H, 4.12; N, 15.61. Found: C, 62.22; H, 4.15; N, 15.52.

**4-Methyl-4-phenyl-3,5-dicyano-2,6-piperidinedione (4j)** : mp 290-291°C; IR (KBr)  $\nu$ : 3304 (NH), 2238 (CN), 1746, 1706 (C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$ : 12.15 (s, 1 H, NH), 7.44-7.20(m, 5 H, ArH), 4.82 (s, 2 H, CH), 1.46 (s, 3 H, CH<sub>3</sub>); Anal. Calcd for C<sub>14</sub>H<sub>11</sub>N<sub>3</sub>O<sub>2</sub>: C, 66.40; H, 4.38; N, 16.59. Found: C, 66.20; H, 4.27; N, 16.40.

## ACKNOWLEDGEMENTS

We are pleased to acknowledge the financial support from Xinxiang Medical College (No. 04GXLP05).

## REFERENCES AND NOTES

1. S. A. El Batran, A. E. N. Osman, M. M. Ismail, and A. M. El Sayed, *Inflammopharmacology*, 2006, **14**, 62.
2. (a) G. M. Ye, Q. Y. Wu, X. J. Huang, Y. Y. Jiang, H. L. Liao, and S. C. Yu, *Dier Junyi Daxue Xuebao*, 2005, **26**, 223; (b) R.W. Holder, J. P. Daub, W. E. Baker, R. H. Gilbert, and N. A. Graf, *J. Org. Chem.*, 1982, **47**, 1445; (c) R. F. Brown and N. M. Van Gulick, *J. Am. Chem. Soc.*, 1955, **77**, 1089; (d) E. B. Reid and T. E. Gompf, *J. Org. Chem.*, 1953, **18**, 661; (e) Y. Xu, Z. H. Zhu, Z. J. Tong, D. M. Peng, and L. X. Duan, *Zhongguo Yiyao Gongye Zazhi*, 1993, **24**, 49; (f) Z. Y. Peng and

- W. Zhu, C. N. Patent 1 740 161, 2005 (*Chem. Abstr.*, 2006, **145**, 103372); (g) S. M. McElvain and D. H. Clemens, *Organic Syntheses*, 1959, **39**, 54.
3. S. M. McElvain and R. E. Lyle, *J. Am. Chem. Soc.*, 1950, **72**, 384.
  4. Y. N. Zhou, Z. W. Fu, and X. J. Wu, C. N. Patent 101 222 041, 2008 (*Chem. Abstr.*, 2008, **149**, 227841).
  5. J. V. Gonna, H. J. Meurer, G. Nover, T. Peun, D. Schonbohm, and G. Will, *Materials Letters*, 1998, **33**, 321.