

HETEROCYCLES, Vol. 91, No. 8, 2015, pp. 1579 - 1584. © 2015 The Japan Institute of Heterocyclic Chemistry
Received, 5th June, 2015, Accepted, 21st July, 2015, Published online, 23rd July, 2015
DOI: 10.3987/COM-15-13261

SITE-SELECTIVE INTRODUCTION OF AN ENAMIDO GROUP AT THE C(3)-POSITION OF INDOLES

Tomoya Miura,* Qiang Zhao, Yuuta Funakoshi, and Masahiro Murakami*

Department of Synthetic Chemistry and Biological Chemistry, Kyoto University,
Katsura, Kyoto 615-8510, Japan

E-mail: tmiura@sbchem.kyoto-u.ac.jp, murakami@sbchem.kyoto-u.ac.jp

Abstract – An enamido group is introduced site-selectively at the C(3)-position of indoles by the rhodium(II)-catalyzed reaction with *N*-sulfonyl-1,2,3-triazoles. Formally, an α -imino rhodium carbene complex is inserted into the C(3)-H bond of an indole.

Indoles are privileged structural motifs found in a myriad of natural products and pharmaceuticals.¹ Numerous methods have been developed for the functionalization of indole skeletons.² Among them, a C-H bond functionalization using metal carbene complexes has received much attention in recent years.³ On the other hand, *N*-sulfonyl-1,2,3-triazoles have emerged as convenient precursors for the generation of α -imino metal carbene complexes.⁴ The complexes contain an electrophilic carbene carbon and a nucleophilic imino nitrogen in the molecule. They can trigger a variety of synthetically useful transformations, including not only typical carbene reactions such as cyclopropanation⁵ and X-H bond insertion (X = carbon,⁶ oxygen,⁷ nitrogen⁸) but also carbene-induced reactions such as [3+2] annulation,⁹ ring expansion,¹⁰ and others.¹¹ Davies and co-workers have demonstrated an excellent method for the functionalization of indole skeletons using α -imino rhodium carbene complexes generated from *N*-sulfonyl-1,2,3-triazoles; 1,3-disubstituted indoles undergo [3+2] annulation reaction in an enantioselective manner to afford pyrroloindolines (Figure 1(a)).^{12,13} This fascinating result led us to





Figure 1. Rh(II)-catalyzed reaction of indoles with *N*-sulfonyl-1,2,3-triazoles


examine the rhodium(II)-catalyzed reaction of indole derivatives having C(3)-H bonds. Now, we report a rhodium(II)-catalyzed reaction of 1,2-disubstituted indoles and 1-monosubstituted indoles with *N*-sulfonyl-1,2,3-triazoles. The presence of the C(3)-H bond shifts the [3+2] annulation pathway to 1,3-insertion of α -imino rhodium carbene complexes into the C(3)-H bond, leading to the formation of β -(3-indolyl)sulfonylenamides with the high (*Z*)-selectivities (Figure 1(b)).

Initially, 4-phenyl-1-tosyl-1,2,3-triazole (**1a**) was prepared from phenylacetylene and tosyl azide according to the authentic procedure using a copper(I) catalyst.¹⁴ When the triazole **1a** (0.2 mmol) was treated with 1,2-dimethyl-1*H*-indole (**2a**, 0.3 mmol) in the presence of (^tBuCO₂)₄Rh₂ (1 mol%) in chloroform (2 mL) at 80 °C, **1a** was completely consumed within 8 hours. Chromatographic purification afforded β -(3-indolyl)sulfonylenamide **3a** in 97% isolated yield with the *Z/E* ratio of >95:5 (Scheme 1). The configuration of the double bond of **3a** was confirmed as *Z* by an NOE study. Unlike the cases with 1,3-disubstituted indoles, no annulation product was formed.¹⁵ We propose the following mechanism for the production of **3a**. α -Diazo imine **1a'** is generated from the triazole **1a** by a ring-chain tautomerization through equilibrium. The transient **1a'** is trapped by a rhodium(II) catalyst to furnish α -imino carbene complex **A**. Nucleophilic addition of the indole **2a** at the C(3)-position to the electrophilic carbene of **A** gives the zwitterionic intermediate **B**. The anionic rhodium of **B** releases an electron pair, which flows onto the imino nitrogen. Since there is a hydrogen atom at the C(3)-position, the imino nitrogen acts as a base to pick up the hydrogen rather than undergoes nucleophilic attack onto the C(2)-carbon atom, furnishing (*Z*)-isomer of **3a**.



Scheme 1. Reaction of 1,2-dimethyl-1*H*-indole (**2a**) with triazole **1a**


1,2-Dimethyl-1*H*-indole (**2a**) was reacted with various triazoles **1** (Table 1). Triazoles **1b–e** possessing aryl and heteroaryl groups at the C(4)-position all reacted well to afford the corresponding products **3b–e** in yields ranging from 90% to 95% with high (*Z*)-selectivities (entries 1–4). The reaction of the alkyl-substituted triazoles **1f–h** gave the products **3f–h**, albeit in moderate yields (entries 5–7).⁷ Aryl group as well as alkyl group were compatible for the R² substituent on the sulfonyl group to afford the products **3i–m** in high yields (entries 8–12).

Table 1. Rh(II)-catalyzed reaction of 1,2-dimethyl-1*H*-indole (**2a**) with various triazoles **1**^a

Entry	1	R ¹	R ²	3	Yield ^b	Z/E ^c	Entry	1	R ¹	R ²	3	Yield ^b	Z/E ^c
1	1b	<i>p</i> -Tol-	<i>p</i> -Tol-	3b	95%	>95:5	7	1h	BzO(CH ₂) ₄ -	<i>p</i> -Tol-	3h	57% ^d	>95:5
2	1c	<i>p</i> -MeO-C ₆ H ₄ -	<i>p</i> -Tol-	3c	96%	>95:5	8	1i	Ph-	<i>p</i> -MeO-C ₆ H ₄ -	3i	98%	>95:5
3	1d	<i>p</i> -CF ₃ -C ₆ H ₄ -	<i>p</i> -Tol-	3d	90%	>95:5	9	1j	Ph-	<i>p</i> -CF ₃ -C ₆ H ₄ -	3j	98%	>95:5
4	1e	3-Thienyl-	<i>p</i> -Tol-	3e	95%	>95:5	10	1k	Ph-	<i>o</i> -Tol-	3k	80%	95:5
5	1f	ⁿ Pr-	<i>p</i> -Tol-	3f	65%	>95:5	11	1l	Ph-	ⁿ Bu-	3l	95%	>95:5
6	1g	^t Bu-	<i>p</i> -Tol-	3g	61% ^d	94:6	12	1m	Ph-	Me-	3m	95%	>95:5

^a Reaction conducted on a 0.2 mmol scale. ^b Isolated yield (average of two runs). ^c The ratio determined by ¹H NMR. ^d **2a** (1.0 mmol, 5.0 equiv) in CHCl₃ (0.5 mL) in the presence of (tBuCO₂)₄Rh₂ (5 mol%).


Next, the use of 1-methyl-1*H*-indole and its derivatives was examined. The C-H bond functionalization took place site-selectively at the C(3)-position, and the corresponding products **3n–s** were obtained in excellent yields (Table 2). In these cases, however, the lower (*Z*)-selectivities (*Z*:*E* = 82:18~88:12) were observed except **3s**, which was ascribed to isomerization of the initially formed (*Z*)-isomer to the (*E*)-isomer occurring during the course of the reaction. In fact, when the reaction of **1a** with 1-methyl-1*H*-indole was monitored by ¹H NMR, the gradual isomerization from (*Z*)-**3n** to (*E*)-**3n** was observed [96:4 (1 h), 83:17 (4 h), 80:20 (8 h)].

Table 2. Rh(II)-catalyzed reaction of various 1-methyl-1*H*-indoles **2** with triazole **1a**^a


3n 97% (83:17) [98% (87:13)] ^b	3o 96% (88:12)	3p 98% (84:16)	3q 96% (82:18)	3r 98% (83:17)	3s 97% (>95:5)

^a The reaction conditions were the same as those in Table 1. Isolated yield (average of two runs). The *Z*:*E* ratio determined by ¹H NMR. ^b In parentheses on a 2.0 mmol scale of **1a**.


Fused indole **2b** with N(1)-to-C(2) was also suitable substrate, giving the product **3t** in 95% yield with the *Z*:*E* ratio of 87:13 (Eq 1).



An all-in-one-pot procedure was also carried out to demonstrate the practical convenience of the present method (Eq 2). Phenylacetylene (**4**), tosyl azide, 1,2-dimethyl-1*H*-indole (**2a**), copper(I) and rhodium(II) catalysts, and chloroform were put in a reaction vessel, and the mixture was stirred at room temperature. After 3 hours, **4** and tosyl azide were both consumed to generate the triazole **1a**. Then, the reaction mixture was stirred at 80 °C for 8 hours, and the following chromatographic purification afforded the product **3a** in 79% yield.



The synthetic utility of the products **3** was exemplified by further transformations. The carbon–carbon double bond of **3a**, **3n**, and **3q** was successfully reduced, giving tryptamine derivatives **5**, **6**, and **7** in good yields when a simple hydrogenation reaction using palladium on charcoal was applied (Eq 3). Furthermore, treatment of **6** with iodine (1.0 equiv) caused electrophilic cyclization to afford fused indole derivative **8** in 77% yield (Eq 4).¹⁶



In summary, a regioselective C-H bond functionalization of 1,2-disubstituted- and 1-monosubstituted indoles by α -imino rhodium carbene complexes is reported. The product selectivity is significantly affected by the substitution pattern at the C(3)-position of the indoles. The resulting products are useful intermediates for the synthesis of tryptamine derivatives.

ACKNOWLEDGEMENTS

This work was supported in part by a Grant-in-Aid for Scientific Research (S) from MEXT and the ACT-C program of the JST. We are grateful to the JSPS for Young Scientists for a Research Fellowship to Y. F. We also thank Dr. Y. Nagata (Kyoto University) for kind help in an X-ray analysis.

REFERENCES AND NOTES

1. Reviews: A. J. Kochanowska-Karamyan and M. T. Hamann, *Chem. Rev.*, 2010, **110**, 4489; P. M. Dewick, 'Medicinal Natural Products: A Biosynthetic Approach, 3rd Ed.', John Wiley & Sons Inc., Chichester, 2009, pp. 311-420.
2. Reviews: S. Cacchi and G. Fabrizi, *Chem. Rev.*, 2005, **105**, 2873; G. Bartoli, G. Bencivenni, and R. Dalpozzo, *Chem. Soc. Rev.*, 2010, **39**, 4449; M. Bandini and A. Eichholzer, *Angew. Chem. Int. Ed.*, 2009, **48**, 9608; C. C. J. Loh and D. Enders, *Angew. Chem. Int. Ed.*, 2012, **51**, 46.
3. Y. Lian and H. M. L. Davies, *J. Am. Chem. Soc.*, 2010, **132**, 440; A. DeAngelis, V. W. Shurtleff, O. Dmitrenko, and J. M. Fox, *J. Am. Chem. Soc.*, 2011, **133**, 1650; T. Goto, Y. Natori, K. Takeda, H. Nambu, and S. Hashimoto, *Tetrahedron: Asymmetry*, 2011, **22**, 907; Y. Cai, S.-F. Zhu, G.-P. Wang, and Q.-L. Zhou, *Adv. Synth. Catal.*, 2011, **353**, 2939; H. Qiu, M. Li, L.-Q. Jiang, F.-P. Lv, L. Zan, C.-W. Zhai, M. P. Doyle, and W.-H. Hu, *Nature Chem.*, 2012, **4**, 733; M. Delgado-Rebollo, A. Prieto, and P. J. Pérez, *ChemCatChem*, 2014, **6**, 2047; L. Li, C. Shu, B. Zhou, Y.-F. Yu, X.-Y. Xiao, and L.-W. Ye, *Chem. Sci.*, 2014, **5**, 4057; L.-Y. Mei, X.-Y. Tang, and M. Shi, *Org. Biomol. Chem.*, 2014, **12**, 1149; Y. Xing, G. Sheng, J. Wang, P. Lu, and Y. Wang, *Org. Lett.*, 2014, **16**, 1244.
4. Reviews: A. V. Gulevich and V. Gevorgyan, *Angew. Chem. Int. Ed.*, 2013, **52**, 1371; H. M. L. Davies and J. S. Alford, *Chem. Soc. Rev.*, 2014, **43**, 5151; P. Anbarasan, D. Yadagiri, and S. Rajasekar, *Synthesis*, 2014, **46**, 3004.
5. S. Chuprakov, S. W. Kwok, L. Zhang, L. Lercher, and V. V. Fokin, *J. Am. Chem. Soc.*, 2009, **131**, 18034; T. Miura, T. Nakamuro, C.-J. Liang, and M. Murakami, *J. Am. Chem. Soc.*, 2014, **136**, 15905.
6. S. Chuprakov, J. A. Malik, M. Zibinsky, and V. V. Fokin, *J. Am. Chem. Soc.*, 2011, **133**, 10352; D. Yadagiri and P. Anbarasan, *Org. Lett.*, 2014, **16**, 2510; S. Park, W.-S. Yong, S. Kim, and P. H. Lee, *Org. Lett.*, 2014, **16**, 4468.
7. T. Miura, T. Biyajima, T. Fujii, and M. Murakami, *J. Am. Chem. Soc.*, 2012, **134**, 194; T. Miura, T. Tanaka, T. Biyajima, A. Yada, and M. Murakami, *Angew. Chem. Int. Ed.*, 2013, **52**, 3883.
8. S. Chuprakov, B. T. Worrell, N. Selander, R. K. Sit, and V. V. Fokin, *J. Am. Chem. Soc.*, 2014, **136**, 195; D. J. Lee and E. J. Yoo, *Org. Lett.*, 2015, **17**, 1830.
9. T. Horneff, S. Chuprakov, N. Chernyak, V. Gevorgyan, and V. V. Fokin, *J. Am. Chem. Soc.*, 2008, **130**, 14972; T. Miura, M. Yamauchi, and M. Murakami, *Chem. Commun.*, 2009, 1470; B. Chattopadhyay and V. Gevorgyan, *Org. Lett.*, 2011, **13**, 3746; M. Zibinsky and V. V. Fokin, *Angew. Chem. Int. Ed.*, 2013, **52**, 1507; S. Chuprakov, S. W. Kwok, and V. V. Fokin, *J. Am. Chem. Soc.*, 2013, **135**, 4652; T. Miura, Y. Funakoshi, and M. Murakami, *J. Am. Chem. Soc.*, 2014, **136**, 2272.
10. T. Miura, Y. Funakoshi, M. Morimoto, T. Biyajima, and M. Murakami, *J. Am. Chem. Soc.*, 2012,

- 134**, 17440; N. Selander, B. T. Worrell, and V. V. Fokin, *Angew. Chem. Int. Ed.*, 2012, **51**, 13054; R. Liu, M. Zhang, G. Winston-McPherson, and W. Tang, *Chem. Commun.*, 2013, **49**, 4376; F. Medina, C. Besnard, and J. Lacour, *Org. Lett.*, 2014, **16**, 3232.
11. E. E. Schultz and R. Sarpong, *J. Am. Chem. Soc.*, 2013, **135**, 4696; B. T. Parr, S. A. Green, and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 4716; J. S. Alford, J. E. Spangler, and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 11712; T. Miura, T. Tanaka, K. Hiraga, S. G. Stewart, and M. Murakami, *J. Am. Chem. Soc.*, 2013, **135**, 13652; J.-M. Yang, C.-Z. Zhu, X.-Y. Tang, and M. Shi, *Angew. Chem. Int. Ed.*, 2014, **53**, 5142; H. Shang, Y. Wang, Y. Tian, J. Feng, and Y. Tang, *Angew. Chem. Int. Ed.*, 2014, **53**, 5662; K. Chen, Z.-Z. Zhu, Y.-S. Zhang, X.-Y. Tang, and M. Shi, *Angew. Chem. Int. Ed.*, 2014, **53**, 6645; E. E. Schultz, V. N. G. Lindsay, and R. Sarpong, *Angew. Chem. Int. Ed.*, 2014, **53**, 9904; D. J. Lee, H. S. Han, J. Shin, and E. J. Yoo, *J. Am. Chem. Soc.*, 2014, **136**, 11606; D. J. Jung, H. J. Jeon, J. H. Kim, Y. Kim, and S. Lee, *Org. Lett.*, 2014, **16**, 2208; A. Boyer, *Org. Lett.*, 2014, **16**, 5878; J. Fu, H. Shen, Y. Chang, C. Li, J. Gong, and Z. Yang, *Chem. Eur. J.*, 2014, **20**, 12881; H.-D. Xu, Z.-H. Jia, K. Xu, H. Zhou, and M.-H. Shen, *Org. Lett.*, 2015, **17**, 66.
12. J. E. Spangler and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 6802.
13. For a thermal reaction of 4-alkoxy-1,2,3-triazoles with indoles, see: J. S. Alford and H. M. L. Davies, *J. Am. Chem. Soc.*, 2014, **136**, 10266.
14. J. Raushel and V. V. Fokin, *Org. Lett.*, 2010, **12**, 4952.
15. The reaction of **1a** with **2a** was carried out under the Davies's conditions (1 mol% (*S*-PTAD)₄Rh₂, cyclohexane, 65 °C, 4 h).¹² However, no annulation product was formed, and **3a** was obtained in 88% yield.
16. Y.-X. Li, H.-X. Wang, S. Ali, X.-F. Xia, and Y.-M. Liang, *Chem. Commun.*, 2012, **48**, 2343.