

HETEROCYCLES, Vol. 98, No. 1, 2019, pp. 88 - 95. © 2019 The Japan Institute of Heterocyclic Chemistry
Received, 22nd October, 2018, Accepted, 26th December, 2018, Published online, 24th January, 2019
DOI: 10.3987/COM-18-14006

REACTION OF 2,3,5,6,7,8-HEXAHYDROQUINOXALINE WITH THIOUREA DERIVATIVES: ANALYSIS OF THE STRUCTURE AND CONFORMATIONAL STATE OF [4.4.3]PROPELLANE

Kazuhide Nakahara,^{a*} Koki Yamaguchi,^b and Hisao Kansui^c

^a Department of Integrative Pharmaceutical Sciences, Faculty of Pharmaceutical Sciences, Setsunan University, 45-1 Nagaotoge-cho, Hirakata, Osaka 573-0101, Japan. E-mail address: kazuhide.nakahara@pharm.setsunan.ac.jp ^b Laboratory of Molecular Design, Faculty of Pharmaceutical Sciences, Sojo University, 4-22-1 Ikeda, Nishi-ku, Kumamoto 860-0082, Japan. ^c Laboratory of Organic Chemistry, Faculty of Pharmaceutical Sciences, Sojo University, 4-22-1 Ikeda, Nishi-ku, Kumamoto 860-0082, Japan

Abstract – The reaction of 2,3,5,6,7,8-hexahydroquinoxaline with thiourea derivatives yields a propellane-type tetraazaindene structure, whose conformation was established by 2D NMR analysis and molecular orbital calculations, and verified by single-crystal X-ray diffraction analysis. Further, the result obtained from X-ray analysis of this propellane-type tetraazaindene structure indicated that the crystalline state was stabilized by specific hydrogen bonds; four hydrogen bonds were of the type $>C=S\cdots H-N$, four of $N-H\cdots N$ type between the imidazolidine moieties, and two of $>C=S\cdots H-C$ type.

Propellane-type structures¹⁻³ possess three rings sharing one C-C covalent bond. Owing to their characteristic stereochemistry, their structural properties, such as bond distances and bond angles, have been well studied.^{4,5} A natural product containing an aza-propellane core, actumine has been isolated from *Sinomenium acutum*⁶ and found to exhibit selective T cell cytotoxicity,⁷ anti-amnesic activity,⁸ and anti-HBV (hepatitis type B virus) activity.⁹ An efficient total synthesis of actumine has also been reported.¹⁰

In a previous study,¹¹ the reaction of dihydropyrazines (**1a**, **1b**, and **2a**),¹² having DNA strand-breakage activity,¹³ with thiourea/methylthiourea provided tetraazaindene derivatives (TAIs) (**3a**, **3b**, **3c**, **3d**, **4a**, and **4b**; Figure 1). Based on X-ray crystallographic data and molecular orbital (MO) calculations, we proposed that the initial stage of cyclization may be triggered by the pericyclic ene reaction between the

NH=C–SH moiety of isothiourea and the C=N moiety of dihydropyrazine.¹¹ Herein we have investigated the conformation of the tetraazaindene moiety in the propellane-type structure (**4a**) by 2D-NMR spectroscopy, MO calculations,¹⁴⁻¹⁷ and X-ray crystallographic analysis.¹⁸⁻²⁰




Figure 1. Structures of compounds **1-4**

The reaction of 2,3,5,6,7,8-hexahydroquinoxaline (**2a**) with thiourea yielded the tetraazaindene propellane-type compound **4a**, whose structure was established by NMR spectroscopy, ¹H-¹H correlation spectroscopy (COSY), and heteronuclear multiple quantum coherence (HMQC) and heteronuclear multiple bond connectivity (HMBC) spectroscopic techniques. Collectively, these techniques indicate the presence of a symmetrical propellane structure in **4a**. Figure 2 shows that the conformation of **4a** was successfully confirmed by the following nuclear Overhauser effect spectroscopic (NOESY) correlations: H-8b and H-11b (1.61 ppm)/H-5a and H-6a (2.52 ppm), H-9b and H-10b (1.40 ppm)/NH-3 and NH-1 (8.01 ppm), and H-5b and H-6b (2.65 ppm)/NH-3 and NH-1 (8.01 ppm). Therefore, the predicted structure of **4a** indicates that both of its six-membered rings adopted boat conformations. Given the long NOESY correlation: H-8b and H-11b/H-5a and H-6a, we assumed that boat conformations were strain. To further confirm the structure of **4a**, MO calculations and single-crystal X-ray diffraction analysis were performed.




Figure 2. Nuclear Overhauser effect (NOE) correlation for predicting the conformation of **4a**

Thus, the NMR results indicated the formation of a symmetrical propellane structure. Figure 3 shows six possible conformations of **4a**. Geometry optimization of these conformations was conducted using density functional theory (DFT) at the B3LYP/6-31G(d) level of theory.^{14,15} By comparing the heats of formation of these conformations, conformation **A** was predicted to be the most stable among the six possible conformations (Figure 3).




Figure 3. Calculated heats of formation (B3LYP/6-31G(d), $^*\Delta\Delta H_f$ (kcal/mol)) of six possible conformations of **4a**. $^*\Delta\Delta H_f$ (kcal/mol) = (conformation **A** – predicted conformations) \times 627.5; 1 hartree = 627.5 kcal/mol

Based on the NMR analysis and MO calculations, we propose that the hydrogen atoms of NH-4 and NH-7 interact with the sulfur atom of the thiocarbonyl group. To investigate this issue in greater detail, the conformations and steric features of **4a** were examined by single-crystal X-ray diffraction analysis.¹⁸⁻²⁰ The ORTEP diagram shown in Figure 4 illustrates that **4a** has a propellane-type tetraazaindene structure. Bond distances and angles relating to the C-C covalent bond (C (3a)–C (7a)) sharing three rings of **4a** are listed in Table 2.




Figure 4. ORTEP diagram of 4a

Table 2. Bond distances and angles of 4a

Bond distances (Å)		Bond angle (°)			
C (3a)-N (3)	1.461	N (3)-C (3a)-N (4)	109.65	N (4)-C (3a)-C (8)	108.45
C (3a)-N (4)	1.461	N (3)-C (3a)-C (8)	111.21	N (4)-C (3a)-C (7a)	113.52
C (3a)-C (8)	1.520	N (3)-C (3a)-C (7a)	101.80	C (8)-C (3a)-C (7a)	112.11
C (3a)-C (7a)	1.581	N (1)-C (7a)-N (7)	110.18	N (7)-C (7a)-C (11)	108.43
C (7a)-N (1)	1.458	N (1)-C (7a)-C (11)	110.93	N (7)-C (7a)-C (3a)	113.41
C (7a)-N (7)	1.458	N (1)-C (7a)-C (3a)	102.21	C (11)-C (7a)-C (3a)	111.60
C (7a)-C (11)	1.527				




Figure 5. Weak interactions in the crystal structure of the propellane-type compound 4a

Table 3. Hydrogen bond interactions (S \cdots H–N, N \cdots H–N, and S \cdots H–C) in the crystal structure of the propellane-type compound **4a**

A \cdots H–D	A \cdots D (Å)	\angle A \cdots H–D (°)
S (1) (B) \cdots H–N (3) (A)	3.685	155.31
S (1) (B) \cdots H–N (4) (A)	3.558	159.18
N (3) (A) \cdots H–N (1) (C)	3.015	161.29
N (4) (C) \cdots H–N (2) (A)	2.988	168.22
S (1) (D) \cdots H–C (A)	3.615	130.14
N (4) (A) \cdots H–N (2) (E)	2.988	168.22
N (3) (E) \cdots H–N (1) (A)	3.015	161.29
S (1) (A) \cdots H–N (3) (F)	3.685	155.31
S (1) (A) \cdots H–N (4) (F)	3.558	159.18
S (1) (A) \cdots H–C (G)	3.615	130.14

*Atoms A and D are the hydrogen bond acceptor and donor, respectively.

Figure 5 shows the crystal structure of **4a** and the associated intermolecular interactions from the surrounding molecules, based on the single-crystal X-ray diffraction analysis. Hydrogen bonds with acceptor and donor are categorized distances of 2.2–2.5 Å as “strong, mostly covalent”, 2.5–3.2 Å as “moderate, mostly electrostatic”, 3.2–4.0 Å as “weak, electrostatic”.²¹ Furthermore, when the \angle A \cdots H–D angle is smaller than 90°, it is thought that hydrogen bonding is not operative.^{22,23}

Table 3 shows the intermolecular hydrogen bonds in the **4a** crystal. It is evident that the **4a** molecule forms ten intermolecular hydrogen bonds. Specifically, four hydrogen bonds are of $>$ C=S \cdots H–N, four of N–H \cdots N type between the imidazolidine moieties, and two of $>$ C=S \cdots H–C type. We propose that these hydrogen bonds contribute to the stabilization of the crystalline state of **4a**. To elucidate the stability of the $>$ C=S \cdots H–N-type hydrogen bond interaction in **4a**, geometry optimization of the composite of the adjoining subunits (A) and (B) (Figure 5), containing two $>$ C=S \cdots H–N-type hydrogen bonds, was conducted using DFT at the B3LYP/6-31G(d) level of theory. The composite of the adjoining (A) and (B) units was more stable because its energy was 5.5 kcal/mol lower than twice the heat of formation of one molecule of **4a**.

In this study, the propellane-type compound **4a** was synthesized and its conformation was elucidated both experimentally and theoretically. We demonstrated that the structure of **4a** in its crystalline state was stabilized by intermolecular hydrogen bonds between its theoretically most stable structures. Thus, we conclude that MO calculations can reliably predict the molecular structure of this propellane-type compound. This knowledge would be useful for future studies of similar propellane-type compounds and tetraazaindene derivatives.

EXPERIMENTAL

MATERIALS AND INSTRUMENTS

The reported melting points are uncorrected. ^1H NMR and ^{13}C NMR spectra were obtained on a JNM-ECA 500 (500 MHz) spectrometer using TMS (tetramethylsilane) as the internal standard. Mass spectra were obtained using a JMS-DX303HF instrument. Chemicals for the synthesis of 2,3,5,6,7,8-hexahydroquinoxaline (**2a**) were purchased from Wako Pure Chemical Industries, Ltd.

GENERAL EXPERIMENTAL PROCEDURE

3a,7a-Tricyclo[3.4.4]butanohexahydro-1*H*-imidazo[4,5-*b*]pyrazine-2(3*H*)-thione (**4a**) was synthesized by the reaction of thiourea and 2,3,5,6,7,8-hexahydroquinoxaline (**2a**).¹¹ Single crystals of **4a** were obtained by recrystallization from MeOH-DMSO; **4a** demonstrated good solubility in a small amount of DMSO but poor solubility in MeOH. Therefore, a large amount of MeOH was added to **4a** that was dissolved in a small amount of DMSO. MeOH was then gradually evaporated, which gave crystals of **4a**.

3a,7a-Tricyclo[3.4.4]butanohexahydro-1*H*-imidazo[4,5-*b*]pyrazine-2(3*H*)-thione (4a): colorless prisms. Yield 68.0%. mp 163–165 °C (dec.). ^1H NMR (500 MHz, DMSO) δ : 1.29 (2H, H-9a, and H-10a), 1.38 (2H, H-8a, and H-11a), 1.40 (2H, H-9b, and H-10b), 1.61 (2H, H-8b, and H-11b), 2.05 (2H, br, NH-4, and NH-7), 2.52 (2H, H-5a, and H-6a), 2.65 (2H, H-5b, and H-6b), 8.01 (2H, NH-1, and NH-3). ^{13}C NMR (125 MHz, DMSO) δ : 20.8 (C9 and C10), 32.8 (C8 and C11), 40.1 (C5 and C6), 74.9 (C3a and C7a), 182.7 (C=S). FAB-MS (*m/z*): 213.2 ($\text{M}^+ + 1$). *Anal.* Calcd for $\text{C}_9\text{H}_{16}\text{N}_4\text{S} \cdot 0.5\text{H}_2\text{O}$: C, 48.84; H, 7.74; N, 25.31: Found: C, 48.56; H, 7.60; N, 25.46.

MOLECULAR ORBITAL (MO) CALCULATIONS

Semi-empirical MO calculations were performed through the CS Chem3D Pro interface and WINMOPAC3.5 using MOPAC2002 on an Intel personal computer.^{16,17} Structures optimized using semi-empirical methods (AM1-, PM5-, or PM6-) were used as the starting geometries for the DFT calculations.^{14,15} The energies were corrected using zero-point vibrational energy (scaled by a factor of 0.9804). Calculations incorporating solvent effect in DMSO, which is a solvent for NMR, used the polarizable continuum model (PCM) method included in the GAMESS software package.¹⁵ As a result of the calculation, there was almost no influence of the solvent effect. The data calculated on the B3LYP/6-31G(d) level are available upon request from the author.

SINGLE-CRYSTAL X-RAY ANALYSIS

A colorless prism crystal having approximate dimensions of $0.500 \times 0.500 \times 0.500$ mm was mounted on a glass fiber. All measurements were made on a Rigaku RAXIS RAPID imaging plate area detector with

graphite monochromated Mo-K α radiation. The data were collected at a temperature of 23 \pm 1 °C to a maximum 2 θ value of 54.9°. The structures were solved by direct method (SIR-2011),¹⁸ and hydrogen atoms were placed at the calculation. A full-matrix least-squares technique was used with anisotropic thermal parameters for non-hydrogen atoms and riding model for hydrogen atoms. All the calculations were performed using the Crystal Structure^{19,20} crystallographic software package.

4a: C₉H₁₆N₄S₁, orthorhombic, space group *Pbca*, a = 15.0969 (6), b = 12.9742 (5), c = 10.7906 (5) (Å), V = 2113.56 (15) (Å³), Density (calc.) 1.334, Density (obs.) 1.334 g cm⁻³, Z = 8, R = 0.0438, Unique data used = 2411, Rw = 0.1752, Goodness of Fit = 1.000. The crystallographic data are deposited at CCDC 1859308.

REFERENCES AND NOTES

1. D. L. Yang, L. Fu, J. R. Li, Q. Zhang, and D. X. Shi, *J. Chem. Res.*, 2016, **40**, 341.
2. M. Shin, M. H. Kim, T.-h. Ha, J. Jeon, K.-H. Chung, J. S. Kim, and Y. G. Kim, *Tetrahedron*, 2014, **70**, 1617.
3. S. Tanaka, K. Seguchi, and A. Sera, *Heterocycles*, 1994, **38**, 2581.
4. W. Wu, J. Gu, J. Song, S. Shaik, and P. C. Hiberty, *Angew. Chem. Int. Ed.*, 2009, **48**, 1407.
5. Y. Maki, M. Tanabe, M. Sako, K. Hirota, and K. Harano, *Chem. Lett.*, 1983, **12**, 1093.
6. M. Tomita, Y. Okamoto, T. Kikuchi, K. Osaki, M. Nishikawa, K. Kamiya, Y. Sasaki, K. Matoba, and K. Goto, *Chem. Pharm. Bull.*, 1971, **19**, 770.
7. B.-W. Yu, J.-Y. Chen, Y.-P. Wang, K.-F. Cheng, X.-Y. Li, and G.-W. Qin, *Phytochemistry*, 2002, **61**, 439.
8. G. W. Qin, X. C. Tang, P. Lestage, D. H. Caignard, and P. Renard, Acutumine and acutumine compounds synthesis and use, US 20060167076A1, 2006.
9. P. Cheng, Y.-b. Ma, S.-y. Yao, Q. Zhang, E.-j. Wang, M.-h. Yan, X.-m. Zhang, F.-x. Zhang, and J.-j. Chen, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 5316.
10. F. Li, S. S. Tartakoff, and S. L. Castle, *J. Am. Chem. Soc.*, 2009, **131**, 6674.
11. K. Nakahara, S. Takechi, K. Yamaguchi, K. Harano, and T. Yamaguchi, *Chem. Pharm. Bull.*, 2009, **57**, 1371.
12. T. Ishiguro and M. Matsumura, *Yakugaku Zasshi*, 1958, **78**, 229.
13. H. Maruoka, N. Kashige, F. Miake, and T. Yamaguchi, *Chem. Pharm. Bull.*, 2005, **53**, 1359.
14. Gaussian 98, Revision A.7. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery, Jr., T. Vreven, K. N. Kudin, J. M. Burant, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rega, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts,

- R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dannenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, J. Cioslowski, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkar, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. Wong, C. Gonzalez, and J. A. Pople, Gaussian, Inc., Pittsburgh PA, 1998.
15. GAMESS: M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, M. Dupuis, and J. A. Montgomery, *J. Comput. Chem.*, 1993, **14**, 1347.
 16. AM1 and PM5 calculations were performed using MOPAC2002 ver. 1.00, Fujitsu Ltd., Tokyo, Japan, 2001.
 17. N. Senda, *Idemitsu Technical Report*, 2006, **49**, 106.
 18. SIR2011: M. C. Burla, R. Caliandro, M. Camalli, B. Carrozzini, G. L. Casciarano, C. Giacovazzo, M. Mallamo, A. Mazzone, G. Polidori, and R. Spagna, *J. Appl. Cryst.*, 2012, **45**, 357.
 19. CrystalStructure 4.3: Crystal Structure Analysis Package, Rigaku Corporation (2000-2018), Tokyo 196-8666, Japan.
 20. CRYSTALS Issue 11: J. R. Carruthers, J. S. Rollet, P. W. Betteridge, D. Kinna, L. Pearce, A. Larsen, and E. Gabe, Chemical Crystallography Laboratory, Oxford, U.K., 1999.
 21. G. R. Desiraju and T. Steiner, *The Weak Hydrogen Bond: In Structural Chemistry and Biology*, Oxford University Press, 2001.
 22. G. A. Jeffrey and S. Takagi, *Acc. Chem. Res.*, 1978, **11**, 264.
 23. G. R. Desiraju, *Acc. Chem. Res.*, 1991, **24**, 290.