

Supporting Information

**CHROMONES FROM THE TWIGS OF *CASSIA FISTULA* AND THEIR
ANTI-TOBACCO MOSAIC VIRUS ACTIVITIES**

**Qiu-Fen Hu,¹ Li-Mei Li,¹ Dong-Lai Zhu,² Zhen-Hua Yu,² Jian-Bo Zhan,² Jie Lou,¹
Ye-De Wang,¹ Kun Zhou,¹ Min Zhou,¹ Yin-Ke Li,¹ Xue-Mei Gao^{1,*}**

¹ Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan Minzu University, Kunming 650031, P. R. China; ² Research and Development of Center, China Tobacco Yunnan Industrial Co., Ltd., Kunming 650231, P.R. China;

E-mail: gao_xuemei@hotmail.com

Contents of Supporting Information

No.	Contents:	Pages:
Figure S1-S4	¹³ C NMR, ¹ H NMR, HSQC, and HMBC spectra of 2-(3-hydroxypropanoyl)-8-methoxy-5-(2-oxopropyl)-4 <i>H</i> - chromen-4-one (1)	3-6
Figure S5-S6	¹³ C NMR and ¹ H NMR spectra of 8-hydroxy-2-(3-hydroxypropanoyl)-5-(2-oxopropyl)-4 <i>H</i> - chromen-4-one (2)	7-8
Figure S7-S8	¹³ C NMR and ¹ H NMR spectra of 2-(2-hydroxyethyl)-7-methoxy-5-(2-oxopropyl)-4 <i>H</i> - chromen-4-one (3)	9-10

Figure S1 ^{13}C NMR spectrum of 2-(3-hydroxypropanoyl)-8-methoxy-5-(2-oxopropyl)-4*H*-chromen-4-one (**1**)




Figure S2 ¹H NMR spectrum of 2-(3-hydroxypropanoyl)-8-methoxy-5-(2-oxopropyl)-4H-chromen-4-one (1)




Figure S3 HSQC spectrum of 2-(3-hydroxypropanoyl)-8-methoxy-5-(2-oxopropyl)-4*H*- chromen-4-one (**1**)





Figure S4 HMBC spectrum of 2-(3-hydroxypropanoyl)-8-methoxy-5-(2-oxopropyl)-4*H*-chromen-4-one



(1)

Figure S5 ^{13}C NMR spectrum of 8-hydroxy-2-(3-hydroxypropanoyl)-5-(2-oxopropyl)-4*H*-chromen-4-one (**2**)




Figure S6 ^1H NMR spectrum of 8-hydroxy-2-(3-hydroxypropanoyl)-5-(2-oxopropyl)-4*H*-chromen-4-one (**2**)




Figure S7 ^{13}C NMR spectra of 2-(2-hydroxyethyl)-7-methoxy-5-(2-oxopropyl)-4*H*-chromen-4-one (**3**)




Figure S8 ^1H NMR spectra of 2-(2-hydroxyethyl)-7-methoxy-5-(2-oxopropyl)-4*H*-chromen-4-one (**3**)

