

**A NEW IRIDOID GLUCOSIDE FROM *ANISACANTHUS VIRGULARIS* AND ITS
ANTIAMOEBIIC ACTIVITY**

**Mohamed S. Refaey,^{1*} Reda A. Abdelhameed,² Mohamed A. A.
Orabi,² Ahmed A. Ali,³ Reda Fouad Ahmed Abdelhameed,⁴
Eman A. A. Mousa,⁵ Shinjiro Hamano,⁶ and Koji Yamada⁷**

¹ Department of Pharmacognosy, Faculty of Pharmacy, University of Sadat City, Menofia 32897, Egypt, ² Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Assiut Branch, Assiut 71524, Egypt, ³ Department of Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut 71526, Egypt, ⁴ Department of Pharmacognosy, Faculty of Pharmacy, Suez Canal University, Ismailia 41522, Egypt, ⁵ Department of Medical Parasitology, Faculty of medicine, South Valley University, Qena, Egypt, ⁶ Department of Parasitology, Institute of Tropical Medicine, Nagasaki University, Sakamoto, 1-12-4, Nagasaki 852-8523, Japan, ⁷ Graduate School of Biomedical Sciences, Nagasaki University, Bunkyo-machi 1-14, Nagasaki 852-8521, Japan; E-mail: mohamed.said@fop.usc.edu.eg

Contents of Supporting Information

No.	Contents	Pages
Figure S1-S7	^{13}C , ^1H -NMR, HH-COSY, HSQC, HMBC, NOESY and mass spectra of 7- <i>O-p-trans</i> coumaroyl caryoptoside (1)	3-6

Figure S1: ^{13}C -NMR spectrum of 7-*O-p-trans* coumaroyl caryoptoside (1)




Figure S2: ^1H -NMR spectrum of 7-*O-p-trans* coumaroyl caryoptoside (1)




Figure S3: HH COSY spectrum of 7-*O-p-trans* coumaroyl caryoptoside (**1**)




Figure S4: HSQC spectrum of 7-*O-p-trans* coumaroyl caryoptoside (**1**)




Figure S5: HMBC spectrum of 7-*O-p-trans* coumaroyl caryoptoside (1)




Figure S6: NOESY spectrum of 7-*O-p-trans* coumaroyl caryoptoside (1)




Figure S7: FAB and HR-FAB-MS spectrum of 7-*O-p-trans* coumaroyl caryoptoside (1)

