

SUPPLEMENTARY INFORMATION

NEW BIOACTIVE SESUQUITERPENOID FROM MALAYSIAN SOFT CORAL GENUS *LEMNALIA*

Kazuki Tani,^a Takashi Kamada,^b Chin-Soon Phan,^{a, c} and Charles S. Vairappan^{a*}

^a Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, 88400 Kota Kinabalu, Sabah, Malaysia. ^b Department of Materials and Life Science, Faculty of Science and Technology, Shizuoka Institute of Science and Technology, 2200-2 Toyosawa, Fukuroi, Shizuoka 437-8555, Japan. ^c Department of Pharmacy, National University of Singapore, 18 Science Dr 4, Singapore 117543, Singapore.

E-Mail: csv@ume.edu.my

SUPPLEMENTARY INFORMATION

Figure S1. ¹H NMR spectrum of parathyrsoidin K (**1**) in CDCl₃ (600 MHz).
Figure S2. ¹³C NMR spectrum of parathyrsoidin K (**1**) in CDCl₃ (150 MHz).
Figure S3. HSQC spectrum of parathyrsoidin K (**1**) in CDCl₃.
Figure S4. ¹H-¹H COSY spectrum of parathyrsoidin K (**1**) in CDCl₃.
Figure S5. HMBC spectrum of parathyrsoidin K (**1**) in CDCl₃.
Figure S6. NOESY spectrum of parathyrsoidin K (**1**) in CDCl₃.
Figure S7. HRESIMS spectrum of parathyrsoidin K (**1**).

Figure S1. ^1H NMR spectrum of parathyroidin K (1) in CDCl_3 (600 MHz).




Figure S2. ^{13}C NMR spectrum of parathyroidin K (1) in CDCl_3 (150 MHz).




Figure S3. HSQC spectrum of parathyroidin K (1) in CDCl₃.




Figure S4. ^1H - ^1H COSY spectrum of parathyroidin K (1) in CDCl_3 .




Figure S5. HMBC spectrum of parathyroidin K (1) in CDCl₃.




Figure S6. NOESY spectrum of parathyrsoidin K (1) in CDCl₃.




Figure S7. HRESIMS spectrum of parathyroidin K (1).

