

HETEROCYCLES, Vol. 88, No. 1, 2014, pp. 377 - 386. © 2014 The Japan Institute of Heterocyclic Chemistry
Received, 17th June, 2013, Accepted, 12th July, 2013, Published online, 19th July, 2013
DOI: 10.3987/COM-13-S(S)33

THE ABILITY OF 1-ARYLTRIAZOLE-CONTAINING NUCLEOBASES TO RECOGNIZE A TA BASE PAIR IN TRIPLEX DNA

Yoshiyuki Hari,* Motoi Nakahara, Shin Ijitsu, and Satoshi Obika*

Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka,
Suita, Osaka 565-0871, Japan; E-mail: hari@phs.osaka-u.ac.jp and
obika@phs.osaka-u.ac.jp

Dedicated to Professor Victor Snieckus on the occasion of his 77th birthday

Abstract – Phosphoramidites bearing propargyl and (*N*-propargylcarbamoyl)methyl groups at the C1-position of deoxyribose were synthesized and introduced into oligonucleotides by using an automated DNA synthesizer. Copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition of the oligonucleotides with various aryl azides led to triplex-forming oligonucleotides (TFOs) possessing the corresponding aryltriazole-containing nucleobases. The triplex-forming ability of TFOs with double-stranded DNA (dsDNA) was evaluated through UV-melting experiments, and it was demonstrated that *m*-hydroxy or *m*-ureido derivatives in the (1-aryltriazol-4-yl)methyl nucleobases likely interacted with a TA base pair in dsDNA.

INTRODUCTION

Triplex formation of double-stranded DNA (dsDNA) by an oligonucleotide (TFO: triplex-forming oligonucleotide) is applicable in various dsDNA-targeting technologies. In triplex formation, TFOs consisting of pyrimidine sequences can sequence-selectively and stably recognize dsDNA via Hoogsteen hydrogen bond formation with AT and GC base pairs in dsDNA by T and C in TFO, respectively. Since no natural nucleic acid specifically recognizes a CG or TA base pair in dsDNA, many studies have been conducted to develop artificial nucleic acids capable of recognizing these base pairs.¹⁻⁴ However, targeting of a TA base pair is difficult because access to the 4-carbonyl oxygen in T is sterically hindered

by the 5-methyl group in T. Therefore, reports on the development of artificial nucleic acids recognizing a TA base pair are scarce.²⁻⁴

We considered that facile and efficient syntheses of various derivatives of a nucleobase structure were necessary for implementing detailed and rational designs of nucleobases for CG or TA base pair recognition, and we have used post-elongation modification (PEM) methods, namely, modification methods after oligonucleotide synthesis, to synthesize derivatives.^{2,5,6} We used a copper-catalyzed alkyne-azide cycloaddition (CuAAC) reaction⁷ as a PEM method and evaluated the various derivatives synthesized; 2-(1-*m*-carbonylamino-phenyl-1,2,3-triazol-4-yl)ethyl nucleobases were recently predicted to interact with the A base of a TA base pair (Figure 1a).² In addition, replacement of the ethylene unit by a methyleneoxy unit led to a significant decrease in the binding affinity to a TA base pair. This may be due to the high flexibility of the methyleneoxy unit. Thus, based on unit length and suppression of the unit flexibility, nucleobases possessing a methylene unit would be of interest (Figure 1a). However, Guianvarc'h *et al.* reported the nucleobase S, which is the N-H group in the amide moiety thought to interact with the 4-carbonyl group in T of a TA base pair (Figure 1b).³ Therefore, nucleobases possessing an amide unit may also be effective as a scaffold for screening nucleobases to recognize a TA base pair (Figure 1b).

Based on previous studies, we synthesized TFOs containing various derivatives of two types of nucleobases, as shown in Figure 1. The TA base pair-recognition ability of the nucleobases was examined by UV melting experiments of triplexes formed with dsDNA.





Figure 1. Nucleobases designed based on our previous results (a) and reports by another group (b)


RESULTS AND DISCUSSION

As shown in Scheme 1, phosphoramidites **1** and **2** possessing ethynyl units to be converted into desired nucleobases after oligonucleotide synthesis were synthesized. Reduction of **3**⁸ with DIBAL followed by treatment with Ohira-Bestmann reagent, (MeO)₂POC(N₂)COMe,⁹ afforded **4** in 82% yield over 2 steps, which was phosphitylated to give the desired **1**. For synthesis of **2**, hydrolysis of **3** under alkaline conditions and condensation with propargylamine in the presence of EDC•HCl and DMAP furnished **5** in 89% yield over 2 steps. The desired phosphoramidite **2** was prepared in 90% yield in the same procedure as **1**.



Scheme 1. Synthesis of the desired phosphoramidites **1** and **2**

Introduction of the synthesized phosphoramidites **1** and **2** into oligonucleotides was performed using an automated DNA synthesizer under conditions of general phosphoramidite chemistry; singly-modified oligonucleotides **6** and **7** as substrates of CuAAC reactions for PEM were successfully synthesized. Through the CuAAC reactions of **6** and **7** with aryl azides¹⁰ under optimized conditions [CuSO₄, sodium ascorbate, tris[(1-benzyl-1*H*-1,2,3-triazol-4-yl)methyl]amine (TBTA)¹¹ and azides in 30% DMSO-phosphate buffer (pH 7.0)],¹² TFOs **8a–f** and **9a–f** bearing the corresponding aryltriazole moieties were obtained, respectively. The purity and molecular weight of all oligonucleotides synthesized were confirmed by reversed-phase HPLC and MALDI-TOF-MS, respectively.



The ability of non-natural nucleobases to recognize a TA base pair was evaluated by UV melting experiments of triplexes formed from TFOs **8** and **9** with a dsDNA target, and the results were compared with those of TFO **10** reported in our previous study.² The obtained T_m values and the difference (ΔT_m) in the T_m values of TFOs possessing any substitution (**b–f**) on the benzene ring from those of unsubstituted congeners (**a**) are summarized in Figure 2. TFO **8a** possessing an unsubstituted phenyl group showed a T_m value of 17 °C, which was drastically lower than that (22 °C) of TFO **10a**. The (1-phenyltriazol-4-yl)methyl unit in TFO **8a** may be spatially incompatible with this dsDNA target in triplex formation because these nucleobases in TFOs **8a** and **10a** likely have no positive interaction with a TA base pair. Based on the results of TFOs **8a–c**, the decrease in T_m values was not observed for the *o*-substituent, unlike TFOs **10a–c**. In contrast, *m*-substituents (**8d–f**) generally increased in the binding affinity to a TA base pair, and hydroxy (**8d**) and ureido (**8f**) groups at the *m*-position led to significantly increased T_m values of +6 °C and +5 °C compared with unsubstituted **8a**, respectively. These results suggest that these analogs positively interact with the TA base pair, for example, via formation of a hydrogen bond with A of a TA base pair. In contrast to the significant stabilization by **8d**, **10d** containing the same *m*-hydroxy group did not stabilize the triplex at all, which is of interest, and further investigation involving a computational study will be required to clarify the reason for this observation. For TFO **8** bearing (1-aryltriazol-4-yl)methyl nucleobases, remarkable changes in T_m values were observed, which were thought to be caused by the lower flexibility of the methylene unit, as expected (Figure 1a). However, the T_m value of TFO **9a** possessing an unsubstituted phenyl group was higher than that of **8a**, but nearly the same as that of **10a**. This result implies that the amide N-H in **9a** does not recognize the T

of a TA base pair, unlike the S found by Sun's group³ (Figure 1b). TFOs **9b–c** with *o*-substituents showed slightly decreased affinity to the TA base pair, while **9d–f** with *m*-substituents showed increased T_m values of +2 °C compared with that of **9a**. However, changes of T_m values were globally low, which may have been caused by the high flexibility of the four-atom length-chain, $-\text{CH}_2\text{CONHCH}_2-$.




Figure 2. Summary of T_m and ΔT_m values obtained by UV melting experiments

In conclusion, two phosphoramidites with ethynyl units were synthesized and introduced into the oligonucleotides. Moreover, through CuAAC reactions of the oligonucleotides, the facile synthesis of TFOs bearing 1-aryltriazole-containing nucleobases was fulfilled. UV melting experiments of synthesized TFOs demonstrated that [1-(*m*-hydroxy- or 1-*m*-ureido-phenyl)triazol-4-yl]methyl nucleobases likely interacted with a TA base pair. Based on this finding and our previous results, the appropriate spacer between aryltriazole and deoxyribose moieties is likely of one- or two-atom length and less flexible. In future studies, we will structurally optimize phenyltriazole and spacer moieties to set the functional group at a suitable position to recognize the TA base pair.

EXPERIMENTAL

All moisture-sensitive reactions were carried out in thoroughly dried glassware under a nitrogen atmosphere. ¹H, ¹³C and ³¹P spectra were recorded on a JEOL JNM-AL300 or JEOL JNM-EX400

spectrometer. Chemical shifts are reported in parts per million referenced to tetramethylsilane ($\delta = 0.00$ ppm) for ^1H NMR spectra, CDCl_3 ($\delta = 77.0$ ppm) and CD_3OD ($\delta = 49.0$ ppm) for ^{13}C NMR spectra, and phosphoric acid ($\delta = 0.00$ ppm) for ^{31}P NMR spectra. IR spectra were recorded on a JASCO FT/IR-4200 spectrometer. Specific rotations were recorded on a JASCO P-2200 polarimeter. EI and FAB mass spectra were measured on a JEOL JMS-600 or JEOL JMS-700 mass spectrometer. MALDI-TOF mass spectra were recorded on a Bruker Daltonics Autoflex II TOF/TOF or JEOL JMS-S3000 mass spectrometer. Fuji Silysia silica gel PSQ-60B (0.060 mm) and FL-60D (0.060 mm) were used for flash column chromatography. For HPLC, SHIMADZU LC-10AT_{VP}, SHIMADZU SPD-10A_{VP} and SHIMADZU CTO-10_{VP} instruments were used. EYELA Cute Mixer CM-1000 was used as a shaker.

1,2-Dideoxy-5-O-(4,4'-dimethoxytrityl)-1-(prop-2-ynyl)- β -D-ribofuranose (4)

Under a nitrogen atmosphere, DIBAL (5.1 mL, 4.74 mmol) was added to a solution of **3**⁸ (1.2 g, 2.37 mmol) in anhydrous CH_2Cl_2 (10 mL) at -78 °C, and the mixture was stirred for 0.5 h. After addition of saturated NaHCO_3 aq., the mixture was extracted with CH_2Cl_2 . The organic extracts were washed with brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:2) to give the aldehyde (1.0 g, 92%). This compound was not subjected to further purification and a portion of this was used in the next step. Under a nitrogen atmosphere, $(\text{MeO})_2\text{POC}(\text{N}_2)\text{COMe}$ (250 mg, 1.30 mmol) and K_2CO_3 (450 mg, 3.24 mmol) were added to a solution of aldehyde (500 mg, 1.08 mmol) in anhydrous MeOH (10 mL) at rt. After being stirred for 5 h, the mixture was concentrated under reduced pressure. The residue was extracted with AcOEt. The organic extracts were washed with water and brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 2:1) to give compound **4** (420 mg, 82% for 2 steps) as a colorless oil. $[\alpha]_{\text{D}}^{24} +8.4$ (*c* 1.06, CHCl_3); IR ν_{max} (KBr) 2932, 1607, 1509, 1462, 1444, 1301, 1251, 1177, 1085, 1035 cm^{-1} ; ^1H -NMR (CDCl_3) δ 1.92–2.02 (3H, m), 2.43–2.49 (2H, m), 3.09 (1H, dd, $J = 6.0, 10.0$ Hz), 3.24 (1H, dd, $J = 4.5, 10.0$ Hz), 3.77 (6H, s), 3.93–3.94 (1H, m), 4.29–4.34 (2H, m), 6.81–6.83 (4H, m), 7.22–7.44 (9H, m); ^{13}C -NMR (CDCl_3) δ 25.0, 39.7, 55.1, 64.5, 70.0, 74.4, 76.3, 80.2, 85.9, 86.1, 113.0, 126.7, 127.7, 128.1, 130.0, 136.0, 144.8, 158.4; MS (EI) m/z 458 (M^+ , 100); HRMS (EI) m/z Calcd for $\text{C}_{29}\text{H}_{30}\text{O}_5$: 458.2093. Found 458.2089.

3-O-[2-Cyanoethoxy(diisopropylamino)phosphino]-1,2-dideoxy-5-O-(4,4'-dimethoxytrityl)-1-(prop-2-ynyl)- β -D-ribofuranose (1)

Under a nitrogen atmosphere, 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite (120 μL , 0.539 mmol) was added to a solution of compound **4** (190 mg, 0.414 mmol) and *N,N*-diisopropylethylamine (210 μL , 0.539 mmol) in anhydrous CH_2Cl_2 (2 mL) at 0 °C, and the mixture was stirred at rt for 1.5 h.

After addition of water, the solvent was removed under reduced pressure and the residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 5:1) to give compound **1** (237 mg, 87%) as a colorless syrup. ¹H-NMR (CDCl₃) δ 1.08 (3H, d, *J* = 7.0 Hz), 1.13–1.18 (9H, m), 1.91–1.96 (2H, m), 2.06–2.21 (1H, m), 2.43–2.46 (1H, m), 2.50–2.55 (2H, m), 2.60 (1H, t, *J* = 6.5 Hz), 3.11–3.16 (2H, m), 3.52–3.81 (10H, m), 4.10–4.15 (1H, m), 4.30–4.33 (1H, m), 4.44–4.48 (1H, m), 6.79–6.83 (4H, m), 7.19–7.36 (7H, m), 7.44–7.47 (2H, m); ³¹P-NMR (CDCl₃) δ 147.1, 147.7; MS (FAB) *m/z* 681 [M+Na]⁺; HRMS (FAB) *m/z* Calcd for C₃₈H₄₇N₂O₆P [M+Na]⁺: 681.3069. Found 681.3093.

2-[1,2-Dideoxy-5-*O*-(4,4'-dimethoxytrityl)-β-D-ribofuranos-1-yl]-*N*-(prop-2-ynyl)ethanamide (5)

KOH (220 mg, 3.95 mmol) was added to a solution of **3**⁸ (1.0 g, 1.97 mmol) in THF/H₂O (1:1, 10 mL) at 0 °C and the mixture was stirred at rt for 12 h. After addition of 2N HCl aq., the mixture was extracted with AcOEt. The organic extracts were washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:6) to give carboxylic acid (970 mg, 96%). This compound was not subjected to further purification and a portion of this was used in the next step. Under a nitrogen atmosphere, propargylamine (27 μL, 0.426 mmol), EDC·HCl (82 mg, 0.426 mmol) and DMAP (9 mg, 0.0711 mmol) were added to a solution of carboxylic acid (170 mg, 0.355 mmol) in anhydrous CH₂Cl₂ (3 mL) at rt and the mixture was stirred for 11 h. After addition of water, the mixture was extracted with CH₂Cl₂. The organic extracts were washed with water and brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:3) to give compound **5** (170 mg, 89% for 2 steps) as a colorless syrup. [α]_D²² -13.8 (*c* 1.12, CHCl₃); IR *n*_{max} (KBr) 3288, 2931, 1656, 1607, 1509, 1444, 1300, 1251, 1082, 1034 cm⁻¹; ¹H-NMR (CDCl₃) δ 1.72 (1H, d, *J* = 3.5 Hz), 1.83 (1H, ddd, *J* = 5.5, 9.5, 13.0 Hz), 2.00 (1H, ddd, *J* = 2.0, 4.0, 13.0 Hz), 2.13 (1H, t, *J* = 2.5 Hz), 2.44 (1H, dd, *J* = 8.5, 15.5 Hz), 2.55 (1H, dd, *J* = 3.0, 15.5 Hz), 3.16 (1H, dd, *J* = 5.5, 10.0 Hz), 3.24 (1H, dd, *J* = 5.0, 10.0 Hz), 3.80 (6H, s), 3.98 (1H, ddd, *J* = 2.5, 5.0, 5.0 Hz), 4.31–4.36 (1H, m), 4.41–4.51 (1H, m), 6.70 (1H, brs), 6.81–6.86 (4H, m), 7.22–7.44 (9H, m); ¹³C-NMR (CDCl₃) δ 28.9, 40.7, 42.0, 55.2, 64.3, 71.3, 73.8, 74.7, 79.6, 86.2, 86.4, 113.1, 126.8, 127.8, 128.1, 130.0, 135.8, 144.7, 158.5, 170.5; MS (EI) *m/z* 515 (M⁺, 100); HRMS (EI) *m/z* Calcd for C₃₁H₃₃NO₆: 515.2308. Found 515.2307.

2-{3-*O*-[2-Cyanoethoxy(diisopropylamino)phosphino]-1,2-dideoxy-5-*O*-(4,4'-dimethoxytrityl)-β-D-ribofuranos-1-yl]-*N*-(prop-2-ynyl)ethanamide (2)

Under a nitrogen atmosphere, 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite (52 μL, 0.233 mmol) was added to a solution of compound **5** (100 mg, 0.194 mmol) and *N,N*-diisopropylethylamine (100 μL, 0.582 mmol) in anhydrous CH₂Cl₂ (2 mL) at 0 °C, and the mixture was stirred at rt for 3 h. After addition of saturated NaHCO₃ aq., the solvent was removed under reduced pressure and the residue was purified

by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:1) to give compound **2** (125 mg, 90%) as a colorless syrup. ¹H NMR (CDCl₃) δ 1.08 (4H, d, *J* = 7.0 Hz), 1.12–1.18 (8H, m), 1.75–1.84 (1H, m), 2.05–2.21 (2H, m), 2.40–2.48 (2H, m), 2.56–2.60 (2H, m), 3.14–3.20 (2H, m), 3.51–3.88 (10H, m), 3.95–4.00 (2H, m), 4.10–4.13 (1H, m), 4.40–4.45 (2H, m), 6.68 (1H, brs), 6.81–6.85 (4H, m), 7.20–7.34 (7H, m), 7.42–7.45 (2H, m); ³¹P-NMR (CDCl₃) δ 147.7, 148.2; HRMS (MALDI-TOF) *m/z* Calcd for C₄₀H₅₀N₃NaO₇P [M+Na]⁺: 738.3279. Found 738.3297.

Synthesis of oligonucleotides

The syntheses of **6** and **7** were performed on a 0.2-μmol scale or 1.0-μmol scale on an automated DNA synthesizer (Gene Design nS-8) using the common phosphoramidite protocol. TFOs synthesized on DMTr-ON mode were cleaved from the CPG resin by treatment with 28% NH₃ aq. at rt for 1.5 h and all the protecting groups on TFOs were removed by treatment with 28% NH₃ aq. at 55 °C for 12 h. The obtained crude TFOs were purified on Sep-Pak[®] Plus C18 cartridges (Waters) followed by reversed-phase HPLC (Waters XBridge[®] OST C18 column 2.5 μm, 10 mm × 50 mm). The composition of the TFOs was confirmed by MALDI-TOF-MS analysis. MALDI-TOF-MS data ([M-H]⁻) for **6** and **7**: **6**, found 4409.26 (calcd. 4409.93). **7**, found 4467.01 (calcd. 4466.99).

Click chemistry: General procedure

A solution of azide compound (5 mM in DMSO, 8 μL) was added to a mixture of CuSO₄ (2 mM in H₂O, 4 μL), TBTA (2 mM in DMSO, 4 μL), sodium ascorbate (10 mM in H₂O, 4 μL), **6** or **7** [0.2 mM in phosphate buffer (pH 7.0), 20 μL] in a 1.5-mL Eppendorf tube. The mixture was shaken at rt using a shaker (1000 rpm) until the reaction was complete. The entire product was purified by reversed-phase HPLC [column: Waters XBridge[®] OST C18 column 2.5 μm, 4.6 mm × 50 mm; eluent: gradient system of MeCN/0.1 M triethylammonium acetate buffer (pH 7.0); flow rate: 1.0 mL/min] to give the desired TFO **8** or **9**. Yields and MALDI-TOF-MS data ([M-H]⁻) for TFOs **8a-f** and **9a-f**: **8a**, 77% yield, found 4529.21 (calcd. 4529.06); **8b**, 71% yield, found 4545.93 (calcd. 4545.01); **8c**, 65% yield, found 4571.01 (calcd. 4572.03); **8d**, 78% yield, found 4545.11 (calcd. 4545.01); **8e**, 84% yield, found 4572.12 (calcd. 4572.03); **8f**, 71% yield, found 4588.02 (calcd. 4587.05); **9a**, 75% yield, found 4586.41 (calcd. 4586.11); **9b**, 73% yield, found 4630.72 (calcd. 4630.12); **9c**, 74% yield, found 4628.68 (calcd. 4629.14); **9d**, 81% yield, found 4602.54 (calcd. 4602.11); **9e**, 80% yield, found 4629.49 (calcd. 4629.14); **9f**, 89% yield, found 4644.31 (calcd. 4644.15).

UV-Melting experiments

UV-Melting experiments were carried out on SHIMADZU UV-1650 and SHIMADZU UV-1800 spectrometers equipped with *T_m* analysis accessory. Equimolecular amounts of the target hairpin-loop dsDNA and TFOs were dissolved in 10 mM sodium cacodylate buffer (pH 6.8) containing 100 mM KCl

and 50 mM MgCl₂ to give a final strand concentration of 1.9 μM. The samples were annealed by heating at 100 °C followed by slow cooling to 5 °C. The melting profiles were recorded at 260 nm from 5 °C to 90 °C at a scan rate of 0.5 °C/min.

ACKNOWLEDGEMENTS

This work was supported in part by a Grand-in-Aid for Scientific Research on Innovative Areas (No. 22136006) from the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT), a Grant-in-Aid for JSPS Fellows of Japan Society for the Promotion of Science (JSPS) and the Mochida Memorial Foundation for Medical and Pharmaceutical Research.

REFERENCES AND NOTES

1. Reviews: S. O. Doronina and J.-P. Behr, *Chem. Soc. Rev.*, 1997, 63; I. Luyten and P. Herdewijn, *Eur. J. Med. Chem.*, 1998, **33**, 515; D. M. Gowers and K. R. Fox, *Nucleic Acids Res.*, 1999, **27**, 1569; K. R. Fox, *Curr. Med. Chem.*, 2000, **7**, 17; J. Robles, A. Grandas, E. Pedroso, F. J. Luque, R. Eritja, and M. Orozco, *Curr. Org. Chem.*, 2002, **6**, 1333; M. G. M. Purwanto and K. Weisz, *Curr. Org. Chem.*, 2003, **7**, 427; S. Buchini and C. J. Leumann, *Curr. Opin. Chem. Biol.*, 2003, **7**, 717; K. R. Fox and T. Brown, *Quart. Rev. Biophys.*, 2005, **38**, 311; D. A. Rusling, V. J. Broughton-Head, T. Brown, and K. R. Fox, *Curr. Chem. Biol.*, 2008, **2**, 1; V. Malnuit, M. Duca, and R. Benhida, *Org. Biomol. Chem.*, 2011, **9**, 326; K. R. Fox and T. Brown, *Biochem. Soc. Trans.*, 2011, **39**, 629; Y. Hari, S. Obika, and T. Imanishi, *Eur. J. Org. Chem.*, 2012, 2875.
2. Y. Hari, M. Nakahara, and S. Obika, *Bioorg. Med. Chem.*, 2013, **21**, in press (doi: [10.1016/j.bmc.2013.05.034](https://doi.org/10.1016/j.bmc.2013.05.034)).
3. D. Guianvarc'h, R. Benhida, J.-L. Fourrey, R. Maurisse, and J.-S. Sun, *Chem. Commun.*, 2001, 1814; D. Guianvarc'h, J.-L. Fourrey, R. Maurisse, J.-S. Sun, and R. Benhida, *Bioorg. Med. Chem.*, 2003, **11**, 2751.
4. S. Verma and P. S. Miller, *Bioconjugate Chem.*, 1996, **7**, 600; O. A. Amosova and J. R. Fresco, *Nucleic Acids Res.*, 1999, **27**, 4632; D. Guianvarc'h, J.-L. Fourrey, R. Maurisse, J.-S. Sun, and R. Benhida, *Org. Lett.*, 2002, **4**, 4209; Y. Wang, D. A. Rusling, V. E. C. Powers, O. Lack, S. D. Osborne, K. R. Fox, and T. Brown, *Biochemistry*, 2005, **44**, 5884; S. Buchini and C. J. Leumann, *Eur. J. Org. Chem.*, 2006, 3152; S. Obika, H. Inohara, Y. Hari, and T. Imanishi, *Bioorg. Med. Chem.*, 2008, **16**, 2945; Y. Hari, S. Kashima, H. Inohara, S. Ijitsu, T. Imanishi, and S. Obika, *Tetrahedron*, 2013, **69**, 6381.
5. Y. Hari, M. Nakahara, J. Pang, M. Akabane, T. Kuboyama, and S. Obika, *Bioorg. Med. Chem.*, 2011, **19**, 1162.

6. Y. Hari, M. Akabane, Y. Hatanaka, M. Nakahara, and S. Obika, [Chem. Commun.](#), 2011, **47**, 4424; Y. Hari, M. Akabane, and S. Obika, [Chem. Commun.](#), 2013, **49**, in press (doi: 10.1039/c3cc44030c).
7. Reviews: H. C. Kolb, M. G. Finn, and K. B. Sharpless, [Angew. Chem. Int. Ed.](#), 2001, **40**, 2004; H. C. Kolb and K. B. Sharpless, [Drug Discov. Today](#), 2003, **8**, 1128.
8. J. Hovinen and H. Salo, *J. Chem. Soc., Perkin Trans. 1*, 1997, 3013.
9. S. Ohira, [Synth. Commun.](#), 1989, **19**, 561; S. Müller, B. Liepold, G. J. Roth, and H. J. Bestmann, [Synlett](#), 1996, 521.
10. All azides used in this study were prepared according to the previous reports.^{2,5,13}
11. T. R. Chan, R. Hilgraf, K. B. Sharpless, and V. V. Fokin, [Org. Lett.](#), 2004, **6**, 2853.
12. M. Nakahara, T. Kuboyama, A. Izawa, Y. Hari, T. Imanishi, and S. Obika, [Bioorg. Med. Chem. Lett.](#), 2009, **19**, 3316.
13. H. Tomioka, T. Matsushita, S. Murata, and S. Koseki, *Liebigs Ann.*, 1996, 1971; J. Andersen, U. Madsen, F. Björkling, and X. Liang, [Synlett](#), 2005, 2209.