

# Case Report

## Distal clavicle osteolysis following fixation with a synthetic ligament

Paul M. C. Dearden, Nicholas A. Ferran, Emyr W. Morris

Access this article online

Website:

[www.internationalshoulderjournal.org](http://www.internationalshoulderjournal.org)

DOI:

10.4103/0973-6042.91003

Quick Response Code:



### ABSTRACT

We present a case of distal clavicle osteolysis following treatment of a chronic acromioclavicular joint dislocation with a synthetic ligament. The relevant literature is reviewed and discussed.

**Key words:** Acromioclavicular joint dislocation, osteolysis, surgilig

Department of Trauma and Orthopaedic Surgery, Lincoln County Hospital, Greetwell Road, Lincoln, LN2 5QY Lincolnshire, United Kingdom

Address for correspondence:

Mr. Nicholas A. Ferran,  
23 Aber Road, Leicester, LE2 2BA. UK.  
E-mail: [nferran@uku.co.uk](mailto:nferran@uku.co.uk)

### INTRODUCTION

The synthetic ligament, Nottingham Surgilig (Surgicraft, UK), was introduced in 2001 by the Nottingham shoulder unit as a method of addressing failed acromio-clavicular joint (ACJ) surgery.<sup>[1]</sup> The artificial braided polyester ligament was developed with a purpose designed loop at each end to facilitate insertion and fixation. The synthetic ligament is looped around the coracoid process, threaded itself, then passed through around the posterior aspect of the clavicle and anchored with a cortical screw, a technique which is quick and requires only two specialized instruments to perform.<sup>[2]</sup> The braided polyester material provides strong biocompatible fixation that develops good tissue ingrowth and provides sufficient stability to allow early shoulder mobilization postoperatively.<sup>[3]</sup> The synthetic ligament has gained popularity over the last three years as a primary method of stabilization in Rockwood classification<sup>[3]</sup> types III-V injuries and is also suitable for use in distal clavicle fractures.

Thus far, the synthetic ligament has had few reported complications such as coracoid fracture,<sup>[4]</sup> ligament failure, and loosening of the screw.<sup>[5]</sup> We report a case of apparent distal clavicle osteolysis in a 56-year-old female patient treated with the synthetic ligament for a chronic Rockwood type III injury.

### CASE REPORT


A 56-year-old lady presented to shoulder clinic with a one-year history of pain over her left dominant acromioclavicular joint. Her pain originated following a fall onto her shoulder

having been knocked off of a pushbike during a collision with a motor vehicle. The primary complaint was of pain, making the patient's work as a machinist difficult. She also complained of a visible but subtle deformity of her acromioclavicular joint resulting in problems keeping the strap of her bra in place.

Clinical examination in the outpatient clinic revealed an obvious acromioclavicular joint dislocation with palpable crepitus on reduction, dynamic instability on clinical testing, and painful movements above 90 degrees of forward flexion and abduction. Radiographs confirmed a diagnosis of grade II ACJ dislocation [Figure 1a]. As she presented with a symptomatic shoulder, she was counseled for an elective ACJ reconstruction. This was a chronic ACJ dislocation and the indication for operation was pain with dynamic instability on clinical examination. At operation, she underwent an uncomplicated ACJ reconstruction using the synthetic ligament technique. Bone quality appeared normal intraoperatively and a chronic grade III injury was noted at time of operation in contrast to the preoperative imaging. Immediate postoperative radiographs were entirely satisfactory.

First follow-up appointment at 2 weeks postoperatively demonstrated no obvious clinical concerns, good wound healing, and no signs of infection or of failure of reconstruction. For the first six weeks postoperative, she undertook passive shoulder exercises under the supervision of the physiotherapists with movement restricted to less than 90 degrees of forward elevation and abduction. At six weeks, this was increased to

Please cite this article as: Dearden PM, Ferran NA, Morris EW. Distal clavicle osteolysis following fixation with a synthetic ligament. *Int J Shoulder Surg* 2011;5:101-4.



**Figure 1:** (a) Disruption of the Acromio-clavicular joint, (b) Immediate postoperative radiograph demonstrating Surgilig screw *in situ*, (c) Follow-up radiograph demonstrating osteolysis of the clavicle around the screw with associated soft tissue swelling, (d) Radiograph taken after removal of implant demonstrating no further bone loss

active range of movement exercises. Follow-up appointment at six weeks revealed no clinical problems and her preoperative symptoms had completely settled.

At 3-month follow-up, the patient reported a firm, painful lump that had developed in the region of the screw. On examination, there was no evidence of fluctuance or erythema, but the swelling was tender to touch. Radiographs demonstrated adequate positioning of the screw with no migration but possible evidence of osteolysis of the surrounding bone [Figure 1b]. The ACJ had however remained reduced with the reconstruction appearing to be intact. At this point, hematological and biochemical inflammatory markers were as follows: WCC 4.9, CRP 2.9, and ESR 6.

At 5-month follow-up, pain and swelling continued to be significant and radiographs demonstrated significant osteolysis around the screw [Figure 1c]. It was thought that the patient was at risk of developing periprosthetic fracture and was listed for urgent removal of the synthetic ligament and screw.

The synthetic ligament and screw were removed six months after their insertion. Moderate tissue in-growth was noted. Perioperatively, there was no evidence of macroscopic infection and this was confirmed on cultures of intraoperative samples. Two weeks post removal of implants, wounds had healed with no evidence of infection and the patient reported a significant improvement in her symptoms of pain. Postoperative inflammatory markers were as follows: WCC 7.9, CRP 2.0, and ESR 7.

Two months following removal of the synthetic ligament, the subtle deformity at time of presentation was noted to have recurred and the patient was again complaining of pain and difficulty working overhead. Radiographs demonstrate displacement of the lateral end of clavicle, but no evidence of further progression of osteolysis [Figure 1d]. A widened acromioclavicular distance is also noted.

She is currently on a waiting list for ACJ reconstruction salvage surgery.

## DISCUSSION

In the operative management of the dislocated ACJ, many surgical techniques have been described. Primary excision of the distal end of the clavicle is usually reserved for symptomatic grade II injuries following a trial of conservative management.<sup>[6,7]</sup> It has been suggested however that in chronic ACJ dislocations, if instability is the main complaint, simple excision of the distal end of the clavicle is not appropriate.<sup>[8]</sup>

In most techniques, the primary aim is to restore the anatomical relationship between coracoid process, acromion, and the distal clavicle. Surgical treatment of ACJ injuries is based around the following three basic principles: reduction of the ACJ by rigid fixation of clavicle to the coracoid; reconstruction of the coracoclavicular ligament by sacrifice of the coracoacromial ligament or other autologous tissue transfer; and by reconstruction of the coraco-clavicular using synthetic material.

Reduction of the ACJ and fixation of acromion and/or coracoid to the clavicle was described by Delbet in 1919 using silver wire<sup>[9]</sup> and has since evolved to use implants such as screws,<sup>[10]</sup> Kirschner-wires,<sup>[11,12]</sup> or variations of a distal clavicle hook plate.<sup>[13,14]</sup>

Distal clavicular excision, with coracoclavicular ligament reconstruction, is a widely used surgical technique. In this procedure, the coracoacromial ligament is sacrificed to recreate the stability provided by the damaged coracoclavicular ligament. Various modifications of the original procedure, described by Weaver and Dunn in 1972,<sup>[15]</sup> have been suggested. The stability and strength of the reduction can be augmented with use of a biological<sup>[16]</sup> or synthetic<sup>[17,18]</sup> coracoclavicular sling. Tendon, muscle, or fascial transfer to recreate the injured coracoclavicular ligament have been widely explored.<sup>[19,20]</sup> Techniques involving dynamic muscle transfer have largely fallen out of favor.<sup>[21]</sup>

Coracoclavicular ligament reconstruction using a prosthetic surgical sling have gained great popularity of late due to ease of surgical technique, strength and longevity of surgical implant, and the ability to preserve surrounding anatomical structures giving a better functional result. Materials such as Ethibond<sup>[22]</sup> or polydioxanone suture (PDS) suture,<sup>[23]</sup> Gore-TEX,<sup>[24]</sup> Dacron<sup>[25]</sup> or braided polyester synthetic ligament such as the "Nottingham Surgilig"<sup>[4]</sup> have been explored with varying degrees of success and complication.

Surgical polyester products similar to the material used in the synthetic ligament has been used in a number of surgical procedures such as vascular grafts/patches,<sup>[26]</sup> mesh in hernia repair,<sup>[27]</sup> and in synthetic reconstruction of anterior cruciate ligament.<sup>[28]</sup>

Reports of foreign body reaction leading to osteolysis became common following the use of polyester synthetic graft material in the reconstruction of anterior cruciate ligaments and lead to suggested limitations to their use in normal healthy individuals.<sup>[28]</sup>

To the best of the authors' knowledge, this is the first report of a sterile foreign body reaction to the polyester material in the synthetic ligament causing osteolysis. The case reported demonstrated negative bacteriology on several occasions and low inflammatory markers which, with the clinical examination of the patient as a correlate, suggest a low level of suspicion of an infective cause. The fact that the patient remained symptomatic even after the removal of the surgical implants further supports this. Our patient was a non-smoker and followed postoperative physiotherapy instruction carefully.

This case also highlights that clinical examination is as important as radiographic imaging. In this case, the imaging suggested a grade II injury; however, clinical examination demonstrated dynamic instability of the ACJ in keeping with the patient's symptoms and difficulty with work and overhead activities. Rolf *et al.* have noted in a series of patients treated surgically for ACJ dislocation; 55% of patients with radiographic diagnosis of grade III injuries were found to have grade IV or V injuries at time of surgery.<sup>[8]</sup>

It is unlikely that screw toggle caused the bone loss in this case, because throughout the investigation of this episode, the radiographic evidence demonstrated the screw to be firmly fixed within the distal clavicle, consistent with the intraoperative findings at the explant stage. We postulate that the osteolysis occurred at the clavicle and not at the coracoid because the clavicular cortex was breached by the screw hole allowing osteolysis around the screw.

Similar anecdotal evidence of aseptic foreign body reactions following ACJ reconstruction, causing periosteal reactions or osteolysis, has been reported. So far, these cases have involved the use of PDS coracoclavicular slings,<sup>[29]</sup> Gore-Tex graft,<sup>[30]</sup> and Dacron.<sup>[31]</sup> Osteolysis of the distal clavicle following ACJ dislocation surgery with similar products suggests the need for further evaluation of the products currently in use to ensure that this is not a more common occurrence than previously anticipated.

Salvage procedures for failed ACJ reconstruction have previously included the use of the synthetic ligament, which in this case would obviously not be appropriate. A simple salvage by performing a distal clavicle osteotomy such as the Mumford procedure may be an option.<sup>[32]</sup> In this case, however, the Mumford procedure may not be of benefit due to the already widened acromioclavicular distance. Modifications of the Weaver-Dunn technique, which involve a distal clavicle osteotomy as part of the technique prior to reduction, may be appropriate.


Although synthetic ligaments can provide stable fixation and is not technically demanding, our case demonstrates that osteolysis is a possible complication of unknown incidence. Bone loss due to osteolysis can make salvage challenging.

## REFERENCES

1. Butt HZ, Wallace WA. Nottingham Surgilig™- A novel device for the surgical treatment of acromioclavicular dislocation. *Orth Prod News* 2005. p. 44-6.
2. Surgicraft UK Product information. Available from: <http://www.surgicraft.co.uk/surgilig.asp> [last accessed on 2011 July 27].
3. Rockwood Jr CA, Young DC. Disorders of the acromioclavicular joint. In: Rockwood Jr CA, Matsen FA 3<sup>rd</sup>, editors. *The Shoulder*. Philadelphia: WB Saunders; 1990. p. 413-68.
4. Jeon IH, Dewnany G, Hartley R, Neumann L, Wallace WA. Chronic acromioclavicular separation: The medium term results of coracoclavicular ligament reconstruction using braided polyester prosthetic ligament. *Injury* 2007;38:1247-53.
5. Bhattacharya R, Goodchild L, Rangan A. Acromioclavicular joint reconstruction using the Nottingham Surgilig: A preliminary report. *Acta Orthop Belg* 2008;74:167-72.
6. Accardo NJ. The Gurd-Mumford procedure for acromioclavicular dislocations. *Bull Tulane Univ Med Fac* 1960;20:41-6.
7. Petersson CJ. Resection of the lateral end of the clavicle. A 3 to 30 year follow-up. *Acta Orthop Scand* 1983;54:904-7.
8. Rolf O, Hann von Weyhern A, Ewers A, Boehm TD, Gohlke F. Acromioclavicular dislocation Rockwood III-V: Results of early versus delayed surgical treatment. *Arch Orthop Trauma Surg* 2008;128:1153-7.
9. Delbet P. Method of treatment of fractures with the collaboration of Marchack (leg) and Lamare and Mass (arm). F. Alcan; 1919.
10. Bosworth BM. Acromioclavicular dislocation; End-results of screw suspension treatment. *Ann Surg* 1948;127:98-111.
11. Mayr E, Braun W, Eber W, Rüter A. Treatment of acromioclavicular joint separations. Central Kirschner-wire and PDS-augmentation. *Unfallchirurg* 1999;102:278-86.
12. Phemister DB. The treatment of acromioclavicular joint by open reduction and threaded-wire fixation. *J Bone Joint Surg* 1942;24:166-8.
13. Cîrstoiu C, Rădulescu R, Popescu D, Ene R, Cîrcotă G, Bădiceanu C. Acroplate - A modern solution for the treatment of acromioclavicular joint dislocation. *J Med Life* 2009;2:173-5.
14. Faraj AA, Ketzer B. The use of a hook-plate in the management of acromioclavicular injuries: Report of ten cases. *Acta Orthop Belg* 2001;67:448-51.
15. Weaver JK, Dunn HK. Treatment of acromioclavicular injuries, especially complete acromioclavicular separation. *J Bone Joint Surg Am* 1972;54:1187-94.
16. Luis GE, Yong CK, Singh DA, Sengupta S, Choon DS. Acromioclavicular joint dislocation: A comparative biomechanical study of the palmaris-longus tendon graft reconstruction with other augmentative methods in cadaveric models. *J Orthop Surg Res* 2007;2:22.
17. Kumar S, Penematsa SR, Selvan T. Surgical reconstruction for chronic painful acromioclavicular joint dislocations. *Arch Orthop Trauma Surg* 2007;127:481-4.
18. LaPrade RF, Wickum DJ, Griffith CJ, Ludewig PM. Kinematic evaluation of the modified Weaver-Dunn acromioclavicular joint reconstruction. *Am J Sports Med* 2008;36:2216-21.
19. Jiang C, Wang M, Rong G. Proximally based conjoint tendon transfer for coracoclavicular reconstruction in the treatment of acromioclavicular dislocation. Surgical technique. *J Bone Joint Surg Am* 2008;90 Suppl 2 Pt 2:299-308.
20. Tauber M, Gordon K, Koller H, Fox M, Resch H. Semitendinosus tendon graft versus a modified Weaver-Dunn procedure for acromioclavicular joint reconstruction in chronic cases: A prospective comparative study. *Am J Sports Med* 2009;37:181-90.

21. Simovitch R, Sanders B, Mehmet O, Lavery K, Warner JJ. Acromioclavicular joint injuries: Diagnosis and management. *J Am Acad Orthop Surg* 2009;17:207-19.
22. Huang T, Hsieh P, Huang KC, Huang KC. Suspension suture augmentation for repair of coracoclavicular ligament disruptions. *Clin Orthop Relat Res* 2009;467:2142-8.
23. Clayer M, Slavotinek J, Krishnan J. The results of coracoclavicular slings for acromio-clavicular dislocation. *Aust N Z J Surg.* 1997;67:343-6.
24. Weinstein DM, McCann PD, McIlveen SJ, Flatow EL, Bigliani LU. Surgical treatment of complete acromioclavicular dislocations. *Am J Sports Med* 1990;18:106-7.
25. Verhaven E, DeBoeck H, Haentjens P, Handelberg F, Casteleyn PP, Opdecam P. Surgical treatment of acute type-V acromioclavicular injuries in athletes. *Arch Orthop Trauma Surg* 1993;112:189-92.
26. Chlupáč J, Filová E, Bacáková L. Blood vessel replacement: 50 years of development and tissue engineering paradigms in vascular surgery. *Physiol Res* 2009;58 Suppl 2:S119-39.
27. Alexandre JH, Bouillot JL. Recurrent inguinal hernia: Surgical repair with a sheet of Dacron mesh by the inguinal route. *Eur J Surg* 1996;162:29-33; discussion 35-6.
28. Bartolozzi P, Salvi M, Velluti C. Long-term follow-up of 53 cases of chronic lesion of the anterior cruciate ligament treated with an artificial Dacron Stryker ligament. *Ital J Orthop Traumatol* 1990;16:467-80.
29. Boldin C, Fankhauser F, Ratschek M, Haller-Schober EM. Foreign-body reaction after reconstruction of complete acromioclavicular dislocation using PDS augmentation. *J Shoulder Elbow Surg* 2004;13:99-100.
30. Stewart AM, Ahmad CS. Failure of acromioclavicular reconstruction using Gore-Tex graft due to aseptic foreign-body reaction and clavicle osteolysis: A case report. *J Shoulder Elbow Surg* 2004;13:558-61.
31. Fullerton LR Jr. Recurrent third degree acromioclavicular joint separation after failure of a Dacron ligament prosthesis: A case report. *Am J Sports Med* 1990;18:106-7.
32. Cook FF, Tibone JE. The Mumford procedure in athletes. An objective analysis of function. *Am J Sports Med* 1988;16:97-100.

**Source of Support:** Nil, **Conflict of Interest:** None declared.



#### Announcement

#### Android App



Download  
**Android  
application**

FREE

A free application to browse and search the journal's content is now available for Android based mobiles and devices. The application provides "Table of Contents" of the latest issues, which are stored on the device for future offline browsing. Internet connection is required to access the back issues and search facility. The application is compatible with all the versions of Android. The application can be downloaded from <https://market.android.com/details?id=comm.app.medknow>. For suggestions and comments do write back to us.