# A cross-sectional study on the association between obesity and changes in the dimensions of the lumbar vertebral column

## Sally Ahmed Magzoub, Qurashi Mohamed Ali

Department of Anatomy, Faculty of Medicine, National University, Khartoum, Sudan

## Abstract

Aim: Obesity is considered a major public health problem and is becoming very common among the population. Impacts of obesity include increased load and weight bearing on the vertebral column, possibly causing alterations in its structure to maintain stability and subsequent disabling lower back pain (LBP). The present study aimed at investigating the association of overweight and obesity with alterations in the lumbar vertebral column as well as LBP. Methodology: Thirty female attendants of the clinic were interviewed with the Roland Disability Questionnaire and specific angles were measured on their lateral lumbosacral X-rays. Results: Changes in the lumbar vertebral column in both the control and study groups were found, but the association of obesity was only significant with vertebral body height of L2, L3, and L5 as well as the back pain disability score. Discussion: The results obtained from this study were comparable and in line with what has been found from previous studies in this field, namely an association between obesity and disabling LBP. While changes in the lumbar angles could not exclusively be correlated with a higher body mass index, changes in the lumbar vertebral height were found to be significantly correlated. Conclusion: Obesity in Sudanese females attending the physiotherapy clinic in Ribat National University Hospital has been proven to be associated with disabling back pain and specific radiographic changes in the lumbar vertebral column. Obesity and its risks on the vertebral column is an association that leaves room for a wide range of studies that should be conducted on a larger scale for more applicable results.

Key words: Angle of curvature, back pain disability score, lumbosacral angle, obesity, vertebral body height

## **INTRODUCTION**

Obesity is becoming a very common and serious problem. According to the WHO fact sheet, updated in 2014, worldwide obesity has doubled since 1980, with >1.4 billion

#### Address for correspondence:

Dr. Sally Ahmed Magzoub, Department of Anatomy, Faculty of Medicine, National University, Khartoum, Sudan. E-mail: barakasally@gmail.com

Access this article online			
Quick Response Code:			
	Website: www.sudanmedicalmonitor.org		
	<b>DOI:</b> 10.4103/1858-5000.185228		

adults aged 20 and above being obese in 2008. Adolescents and children are also included in this pandemic, i.e., growing worldwide, in both developed and developing countries.<sup>[1-3]</sup> In this modern age, a combination of affordable high-calorie foods and sedentary lifestyle can be implicated as one of the main causes of these rising statistics. Obesity, which is a preventable condition, is defined as an abnormal or excessive accumulation of body fat in relation to the individuals' height that may impair health.<sup>[3]</sup>

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Magzoub SA, Ali QM. A cross-sectional study on the association between obesity and changes in the dimensions of the lumbar vertebral column. Sudan Med Monit 2016;11:35-40.

Various parameters are used to give obesity a reliable numerical value;<sup>[4]</sup> the most frequently used parameter being the body mass index (BMI) calculated as the person's weight in kg divided by their height in meters squared. A value  $\geq 25$  is considered overweight, whereas obesity falls in the range of  $\geq 30$  (also classified as Grade 1 if  $\geq 30$ , Grade 2 if ranging from 35 to 40, and Grade 3 if  $\geq 40$ ).<sup>[5]</sup>

Concerns regarding obesity stem from its high association with diseases and physical problems<sup>[6]</sup> as well as its rapid growth rate.<sup>[7]</sup> Excessive weight results in abnormal mechanics which may explain the musculoskeletal disorders obese patients tend to suffer from.<sup>[8,9]</sup> Negative effects of obesity on the vertebral column include poor flexibility, increased stiffness, and weak back muscles. The curvature of the vertebral column increases specifically in the lower back, causing an increase in the lumbar lordosis and pelvic tilt.

About 3.4 billion adults die annually from conditions resulting from obesity. Studies have shown that there is an association between high range BMI's, increased waist circumference, and chronic conditions including hypertension, ischemic heart disease, certain cancers (breast, prostate, bowel) and type 2 diabetes mellitus. Earlier onset of obesity (in childhood and adolescence) is more likely to continue into adulthood which has been linked to a 50–100% higher morbidity from obesity-related diseases.Obesity has been linked to psychiatric disorders such as depression and anxiety.

Obesity also causes decreased bone density and strength (osteoporosis) leading to a higher risk of fractures in the vertebral column. Osteoarthritis damages the joints between the facets thereby decreasing their mobility. It is therefore very common for obese people to suffer from chronic back pain.

Obesity has been linked to spinal impairment including disc degeneration, modified spinal posture increased lumbar lordosis angle, which all increase the chances of lower back pain (LBP).<sup>[10-12]</sup> A vast number of studies have been conducted in relation to this subject and to date, results have either confirmed this association or shown that further research is required to prove that there is one, depending on the aspect of spinal impairment being investigated.

Considering the functional anatomy of the vertebral column, it is arranged such that it has an anterior weight bearing portion and a posterior portion that serves to protect the spinal cord and its associated structures.<sup>[13-15]</sup>

Impairment of the vertebral column as a result of obesity has been studied using a number of techniques including the evaluation of lateral lumbosacral X-rays and magnetic resonance imaging (MRI).<sup>[16]</sup> Specific angles and measurements are taken on these images and correlated with parameters used to define obesity. These include the lumbosacral angle, which is regularly quoted as an important measurement in evaluating the etiology of LBP. The lumbar lordosis angle defines the curvature of the lumbar portion of the vertebral column, which is a significant weight bearing segment. It serves to reduce stiffness and absorb impact. The intervertebral discs (IVDs) also play a major role in stabilizing the vertebral column while and distributing forces during its movement. Damage and effects on the IVD may be reflected as disc degeneration, reduction of specific angles on X-ray or altered signal frequencies on MRI images.<sup>[17,18]</sup>

This study aimed to investigate whether obesity in Sudanese females suffering from LBP was associated with changes in the lumbar vertebral column, testing the hypothesis that a BMI of >30 is more likely to be associated with changes in the lumbar vertebral column.

# **METHODOLOGY**

This study was conducted as a cross-sectional study at the physiotherapy clinic at Ribat National University Hospital in the time period between November 2014 and early January 2015. A total of 30 females with LBP were included in the study; those with a BMI of <25, history of back trauma or spinal surgery were excluded from the study. All subjects were given a verbal explanation of the project and participated in the study voluntarily after giving consent. For scoring of their back pain, each patient was interviewed with the Roland Disability Questionnaire (RDQ) (12 item version).<sup>[19-21]</sup> Following the interview, each patient's weight was taken using a standard bathroom scale and their height was measured as well. They were then requested to take a lateral lumbosacral X-ray, which was used for the measurement of the lumbar dimensions.

Measurements on the lumbosacral X-rays involved the following procedures;

## The lumbosacral angle

A line was drawn following the lower margin of L5 and another one following the upper margin of S5. The two lines were extended until they intersected and the angle between them was measured [Figure 1].

## The angle of curvature

A line was drawn along the upper margin of L1 as well as the lower margin of L5. Both lines were extended to their point of intersection then bisected at 90°. The acute angle formed by the intersection of these bisecting lines was then measured [Figure 2].

## The intervertebral discs angle between L4 and L5

A line was drawn along the lower border of L4 and another one along the upper border of L5. The angle between these two lines was measured [Figure 3].

#### Midline height of the vertebral bodies

Lines were drawn along the upper and lower borders of each vertebral body from L1 to L5. The midpoint of these lines was measured and connected to measure the mid-height of each vertebral bod [Figure 4].

All lines were drawn on the X-rays using an erasable red marker. Each measurement was taken twice and the average recorded on the designated form.

Statistical analysis was done using SPSS (Version 19) (IBM Corp. Released 2010. IBM SPSS Statistics for Windows, Version 19.0. Armonk, NY: IBM Corp). Descriptive analysis included the calculation of the means

and standard deviations for all measurements attained. Association was measured using the independent *t*-test with a significant relationship being a numerical value of 0.05 or less (P < 0.05).

## RESULTS

Thirty female subjects were included in the study, all complaining of LBP for more than a years' duration. The youngest participant was 26-year-old, whereas the eldest was 60 years of age; the average age of the subjects included in the study was 45 years. The control group included any female with a BMI ranging from 25 to 29. This group comprised 10 out of the total 30 females. Their average BMI was 27.3. The remaining 20 were classified as the study group, including females with a BMI of 30 or higher. Their average BMI was 36 [Figure 5].



Figure 1: Measurement of the lumbosacral angle. Source: www.quia. com



Figure 3: Measurement of the intervertebral disc angle



Figure 2: Measurement of the angle of curvature. Source: www. hindawi.com



Figure 4: Measurement of mid vertebral height

Seventy-three percent of the subjects (22 out of the 30) had a lumbosacral disc angle value of less than the normal range (15–20°), whereas 20% of them (6 out of the 30) fell within the normal range. The remaining 7% (2 subjects) had a lumbosacral angle of >20 [Figure 6].

A larger proportion of the subjects (57%, 17 out of the original 30) had an angle of curvature (lumbar lordosis angle) more than the upper limit of normal (45°) indicating hyperlordosis [Figure 7].

Vertebral height averages for all subjects were 2.2 cm (L1), 2.8 cm (L2), 3.0 cm (L3), 2.8 cm (L4), and 2.7 cm (L5) [Figures 8-10].

Considering the disability due to back pain scores, the average score for participants was 8. This is considered severe disability if taken in the context of the RDQ scoring system where a 40–60% score is interpreted as severe<sup>[22]</sup> and the average score in this study was 66%.

## DISCUSSION

Taking advantage of X-ray survey of forward and lateral projections can be sufficient to fully appreciate the bone component parameters of the lumbar spine. In this study, lateral images of the lumbar vertebral column were utilized.



Figure 5: Number of subjects and their body mass index ranges



Figure 7: Lumbar lordosis/angle of curvature

The WHO classifies people with a BMI of 25 as being overweight while those with a BMI of 30 as being obese and as stated earlier, increased weight leads to increased forces acting on the vertebral column requiring functional adjustments.

While changes in the lumbar dimensions were found in both groups [Table 1], statistical analysis of the data indicated that very few of these differences were actually significant enough to prove an association with a higher BMI.

The values for the lumbosacral angle (P = 0.073), angle of curvature (P = 0.4), and IVD angle between L4 and L5 (P = 0.365); all had P > 0.05 and were therefore not considered significantly associated with a BMI of >30. These results are comparable to the work conducted by Romero-Vargas. The parameters investigated in that study were spinopelvic but did include part of the lumbar column; their results indicated a poor correlation similar to the poor association found in this study.

When considering the lumbar vertebral height, a significant P value was found for lumbar vertebrae numbers 2 (P = 0.000) 3 (P = 0.000) and 5 (P = 0.005) [Table 2]. Information from review of the literature did not yield any information regarding the vertebral body height and



Figure 6: Lumbosacral disc angle values



Figure 8: Average mid vertebral body heights



Figure 9: Narrowing of the disc space

Table 1: The results of the measurements obtained for the lumbar parameters of the control (body mass index <30) and study (body mass index >30) groups

Parameter	Control group		Study group	
	Mean	SD	Mean	SD
Lumbosacral angle	10.70	1.567	13.45	4.501
Angle of curvature	45.10	2.726	43.85	4.196
Intervertebral disc angle (L4 and L5)	12.60	0.516	13.95	4.582
Mid vertebral body height				
L1	2.510	0.1912	2.080	0.4720
L2	3.230	0.1252	2.410	0.4278
L3	3.690	0.1912	2.480	0.3901
L4	2.980	0.0632	2.680	0.5126
L5	2.980	0.0632	2.415	0.5788

SD: Standard deviation

Table 2: The results of the statistical analysis obtained to compare the parameters between the control and study group using the independent *t*-test

Parameter	<i>t</i> -test	Significant
Lumbosacral angle	1.862	0.073
Angle of curvature	-0.852	0.4
Intervertebral disc angle (L4 and L5)	0.921	0.365
Mid vertebral body height		
L1	-2.751	0.1
L2	-5.889	0.000
L3	-9.21	0.000
L4	-1.828	0.078
L5	-3.05	0.005

obesity, so this particular aspect was difficult to compare with previous results. However, these results indicate a



Figure 10: Intersection of the measurement lines within the intervertebral foramen

significant association between the vertebral heights of these selected vertebrae and a higher BMI.

Finally, a significant association was also found between a higher BMI and the disability due to back pain score (P = 0.046) consistent with Guo *et al.*<sup>[23]</sup> who, using the same disability score found an increased disability score in obese patients. This finding is further supported by several sources from the literature. Adults with disabilities have a higher rate of obesity that those without disabilities.<sup>[24-27]</sup>

# CONCLUSION

The results of this study are generally in line with what has been found in previous studies, proving that a higher BMI is associated with disabling LBP. The aspect related to changes in the dimensions of the lumbar column indicated that changes, while seen, cannot directly be linked to a higher BMI following comparison of the study and control groups. This may have been due to the small sample size and future research in this area should include a larger sample size as well as BMI ranges that are distinct from each other (for example comparing normal BMI with extremely high BMI groups). Other areas of research regarding this particular topic may be to study the effect of weight loss on the degree of disability and vertebral changes.

#### **Acknowledgments**

The authors would like to acknowledge

- Dr. Asim Qurashi for his assistance with the writing of this article
- Dr. Eman Hashim for contributing to the statistical analysis of the data.

#### **Financial support and sponsorship**

Nil.

#### **Conflicts of interest**

There are no conflicts of interest.

## REFERENCES

- He F, Liu J. Prevalence of obesity among primary students from 2009 to 2014 in China: An update meta-analysis. Int J Clin Exp Med 2014;7:5348-52.
- Rohilla R, Rajput M, Rohilla J, Malik M, Garg D, Verma M. Prevalence and correlates of overweight/obesity among adolescents in an urban city of North India. J Family Med Prim Care 2014;3:404-8.
- Aronne LJ. Classifi cation of obesity and assessment of obesityrelated health risks. Obes Res 2002;10 Suppl 2:105S-15S.
- 4. Borruel S, Moltó JF, Alpañés M, Fernández-Durán E, Álvarez-Blasco F, Luque-Ramírez M, et al. Surrogate markers of visceral adiposity in young adults: Waist circumference and body mass index are more accurate than waist hip ratio, model of adipose distribution and visceral adiposity index. PLoS One 2014;9:e114112.
- Contardo Ayala AM, Nij pels G, Lakerveld J. Validity of self-measured waist circumference in adults at risk of type 2 diabetes and cardiovascular disease. BMC Med 2014;12:170.
- Shazia SM, Badaam KM, Deore DN. Assessment of aerobic capacity in overweight young females: A cross-sectional study. Int J Appl Basic Med Res 2015;5:18-20.
- Gatineau M, Hancock C, Dent M. Adult disability and obesity. Oxford: National Obesity Observatory; 2013.
- Bolgen-Cimen O, Arýncý-Yncel N, Karabiber M, Erdogan C. Role of obesity in low back pain related disability. West Indian Med J 2007;56:252.
- Laslett LL, Just Nee Foley SJ, Quinn SJ, Winzenberg TM, Jones G. Excess body fat is associated with higher risk of vertebral deformities in older women but not in men: A cross-sectional study. Osteoporos Int 2012;23:67-74.
- Liuke M, Solovieva S, Lamminen A, Luoma K, Leino-Arjas P, Luukkonen R, *et al.* Disc degeneration of the lumbar spine in relation to overweight. Int J Obes (Lond) 2005;29:903-8.
- 11. Vismara L, Menegoni F, Zaina F, Galli M, Negrini S, Capodaglio P. Effect of obesity and low back pain on spinal mobility: A cross sectional study in women. J Neuroeng Rehabil 2010;7:3.
- 12. Romero-Vargas S, Zárate-Kalfópulos B, Otero-Cámara E,

Rosales-Olivarez L, Alpízar-Aguirre A, Morales-Hernández E, *et al.* The impact of body mass index and central obesity on the spinopelvic parameters: A correlation study. Eur Spine J 2013;22:878-82.

- Rawls A, Fisher RE. Development and functional anatomy of the spine. In: The Genetics and Development of Scoliosis. 12<sup>th</sup> ed. USA: Springer; 2010.
- Damasceno LH, Catarin SR, Campos AD, Defino HL. Lumbar lordosis: A study of angle values and of vertebral bodies and intervertebral discs role. Acta Ortop Bras 2006;14: 193-8.
- Schafer RC. General spinal biomechanics. In: Clinical Biomechanics: Musculoskeletal Actions and Reactions. 2<sup>nd</sup> ed. Philadelphia, Pennsylvania, United States: Williams and Wilkins; 1987.
- Samartzis D, Karppinen J, Chan D, Luk KD, Cheung KM. The association of lumbar intervertebral disc degeneration on magnetic resonance imaging with body mass index in overweight and obese adults: A population-based study. Arthritis Rheum 2012;64:1488-96.
- Takatalo J, Karppinen J, Taimela S, Niinimäki J, Laitinen J, Sequeiros RB, *et al.* Association of abdominal obesity with lumbar disc degeneration – A magnetic resonance imaging study. PLoS One 2013;8:e56244.
- 18. Ablyazov O. X ray parameters of the lumbar spine. MHSJ 2012;10:37-40.
- Turner JA, Fulton-Kehoe D, Franklin G, Wickizer TM, Wu R. Comparison of the Roland – Morris disability questionnaire and generic health status measures: A population-based study of workers' compensation back injury claimants. Health Serv Res 2003;28:1061-7.
- Stratford PW, Binkley J, Solomon P, Finch E, Gill C, Moreland J. Defining the minimum level of detectable change for the Roland-Morris questionnaire. Phys Ther 1996;76:359-65.
- 21. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity and low back pain: A meta-analysis. Am J Epidemiol 2010;171:135-54.
- Ojoawo AO, Oloagun MO, Bamiwoye SO. Relationship between pain intensity and anthropometric indices in women with low back pain – A cross sectional study. J Phys Ther 2011;3:45-51.
- Guo JM, Zhang GQ, Alimujiang. Effect of BMI and WHR on lumbar lordosis and sacrum slant angle in middle and elderly women. [In Chinese] Zhongguo Gu Shang 2008;21:30-1.
- 24. Styne DM. Childhood and adolescent obesity. Prevalence and significance. Pediatr Clin North Am 2001;48:823-54, vii.
- 25. Aronne LJ. Classification of obesity and assessment of obesityrelated health risks. Obes Res 2002;10 Suppl 2:105S-15S.
- Longo UG, Loppini M, Denaro L, Maff ulli N, Denaro V. Rating scales for low back pain. Br Med Bull 2010;94:81-144.
- 27. Gatineau M, Hancock C, Dent M. Adult disability and obesity. Oxford: National Obesity Observatory; 2013.